

On quasi-Hopf smash products and twisted tensor products of quasialgebras

Helena Albuquerque*

Departamento de Matematica-Faculdade de Ciencias e Tecnologia,
Universidade de Coimbra, Apartado 3008, 3000 Coimbra, Portugal
e-mail: lena@mat.uc.pt

Florin Panaite†

Institute of Mathematics of the Romanian Academy
PO-Box 1-764, RO-014700 Bucharest, Romania
e-mail: Florin.Panaite@imar.ro

Abstract

We analyze some relations between quasi-Hopf smash products and certain twisted tensor products of quasialgebras. Along the way we obtain also some results of independent interest, such as a duality theorem for finite dimensional quasi-Hopf algebras and a universal property for generalized diagonal crossed products.

Introduction

If (A, μ_A, u_A) and (B, μ_B, u_B) are algebras in a monoidal category $\mathcal{C}$ and $R : B \otimes A \rightarrow A \otimes B$ is a morphism in $\mathcal{C}$ satisfying a certain list of axioms, then $A \otimes B$ becomes also an algebra in $\mathcal{C}$, with a multiplication defined in terms of μ_A , μ_B and R . This construction appeared in a number of contexts and under different names. Following [15] we call such an R a *twisting map* and the algebra structure on $A \otimes B$ afforded by it the *twisted tensor product* of A and B and denote it by $A \otimes_R B$ (if A and B are ordinary associative algebras and R is the usual flip, then $A \otimes_R B$ coincides with the usual tensor product of algebras). Analogues of twisting maps for monads and operads are known as *distributive laws*, see for instance [4], [25], [30]. The twisted tensor product of associative algebras can be regarded as a representative for the cartesian product of noncommutative spaces, better suited than the ordinary tensor product, see [15], [18], [20] for a detailed discussion and references. Prominent examples of twisted tensor products of algebras are the so-called *braided tensor products*, which are part of the "braided geometry" developed by Majid in the early 1990's; namely, a braiding on a monoidal category provides a twisting map between any two algebras in the category. Hopf algebra theory provides also plenty of examples of twisted tensor products, in particular the usual smash product $A \# H$ is the twisted tensor product $A \otimes_R H$, with $R : H \otimes A \rightarrow A \otimes H$, $R(h \otimes a) = h_1 \cdot a \otimes h_2$.

*The first author was partially supported by the Centre for Mathematics of the University of Coimbra (CMUC)

†Research carried out while the second author was visiting the Centre for Mathematics of the University of Coimbra (CMUC) supported by a postdoctoral fellowship offered by CMUC. This author was also partially supported by the programme CEEEX of the Romanian Ministry of Education and Research, contract nr. CEx05-D11-11/2005.

The term *quasialgebra* was introduced in [2], to designate an algebra in a monoidal category. We restrict here the term to algebras in monoidal categories associated to a quasi-bialgebra or quasi-Hopf algebra H , such as ${}_H\mathcal{M}$, ${}_H\mathcal{M}_H$, ${}^H_H\mathcal{YD}$. Since these categories have nontrivial associators, such quasialgebras are usually *nonassociative* as algebras, but their lack of associativity is (well) controlled by the associators of the category. Important examples of quasialgebras are the octonions and the other Cayley algebras, see [2]. Another class of examples of interest for us here is obtained as follows: if H is a quasi-Hopf algebra, B an associative algebra and $v : H \rightarrow B$ an algebra map, then on B one can introduce a structure of algebra in ${}_H\mathcal{M}$, denoted by B^v , see [12]. For the particular case $B = H$, $v = id_H$, the quasialgebra B^v is denoted by H_0 , and by [11] it follows that it is actually an algebra in the Yetter-Drinfeld category ${}^H_H\mathcal{YD}$.

A feature of quasialgebras in this sense, not available for algebras in arbitrary monoidal categories (unless very special conditions hold, cf. [29]) is the fact that there exist several *crossed products* associated to them, which are usually *associative* algebras. For instance, if H is a quasi-bialgebra and A is an algebra in ${}_H\mathcal{M}$, one can consider the smash product $A\#H$ introduced in [12], an associative algebra generalizing the classical Hopf smash product and sharing many of its properties. Other examples are diagonal crossed products, L-R-smash products, two-sided smash and crossed products, see [13], [17], [28].

These quasialgebras are also part of an emerging *nonassociative geometry* (cf. [1], [5], [24]), regarded as a further extension of noncommutative geometry, with the “coordinate algebra” allowed to be nonassociative. By analogy with the associative situation, the twisted tensor product of quasialgebras might be regarded as a representative for the cartesian product of “nonassociative spaces”.

The aim of this paper is to study certain classes of twisted tensor products of quasialgebras and their relations with quasi-Hopf smash products. If H is a quasi-Hopf algebra, A a left H -module algebra and C an algebra in the Yetter-Drinfeld category ${}^H_H\mathcal{YD}$, we consider an object denoted by $A \diamond C$, which is a certain twisted tensor product in the category ${}_H\mathcal{M}$ and is defined in such a way that if A is also an algebra in ${}^H_H\mathcal{YD}$ then $A \diamond C$ coincides with the braided tensor product $A \underline{\otimes} C$ in ${}^H_H\mathcal{YD}$. We are mainly interested in the case $C = H_0$, so we have the left H -module algebra $A \diamond H_0$ and we want to see how is it related to the smash product $A\#H$. For Hopf algebras this question has a trivial answer: $A\#H$ and $A \diamond H_0$ coincide. This cannot be the case for proper quasi-Hopf algebras, because $A\#H$ is associative while $A \diamond H_0$ is *not*. The answer is the following: $A \diamond H_0$ is isomorphic as left H -module algebras to $(A\#H)^j$, where $j : H \rightarrow A\#H$, $j(h) = 1\#h$. If A is moreover an algebra in ${}^H_H\mathcal{YD}$, we prove that $(A\#H)^j$ becomes also an algebra in ${}^H_H\mathcal{YD}$ in a natural way and we have $A \underline{\otimes} H_0 \simeq (A\#H)^j$ as algebras in ${}^H_H\mathcal{YD}$.

More can be said if A is H_0 itself. For this, we prove a result of independent interest: an algebra isomorphism $B^v\#H \simeq B \otimes H$, the quasi-Hopf analogue of a well-known result from Hopf algebra theory (see [26]). As a consequence, we obtain $H_0 \underline{\otimes} H_0 \simeq (H \otimes H)^\Delta$ as algebras in ${}^H_H\mathcal{YD}$. As another consequence, also of independent interest, we obtain a duality theorem, stating that if H is a finite dimensional quasi-Hopf algebra then the two-sided crossed product $H \rtimes H^* \ltimes H$ introduced by Hausser and Nill in [17] is isomorphic to $End(H) \otimes H$ as algebras.

In the last two sections we analyze some iterated products (in particular in the sense of [18]) involving quasialgebras, and some universal properties. For instance, we find a relation between the universal property of $A\#H$ and the universal property of $A \diamond H_0$ regarded as a twisted tensor product of quasialgebras, and we find a new kind of universal property of the smash product $A\#H$, obtained as an immediate consequence of a very general universal property for generalized diagonal crossed products (as in [13]), which we obtain inspired by the universal property of the diagonal crossed product $H^* \bowtie \mathbb{A}$, formulated by Hausser and Nill in [17].

1 Preliminaries

In this section we recall some definitions and results and we fix some notation that will be used throughout the paper.

We work over a base field k . All algebras, linear spaces etc. are over k ; unadorned $\otimes$ means $\otimes_k$. Following Drinfeld [16], a quasi-bialgebra is a fourtuple $(H, \Delta, \varepsilon, \Phi)$, where H is an associative algebra with unit 1, Φ is an invertible element in $H \otimes H \otimes H$, and $\Delta : H \rightarrow H \otimes H$ and $\varepsilon : H \rightarrow k$ are algebra homomorphisms satisfying the identities (for all $h \in H$):

$$(id \otimes \Delta)(\Delta(h)) = \Phi(\Delta \otimes id)(\Delta(h))\Phi^{-1}, \quad (1.1)$$

$$(id \otimes \varepsilon)(\Delta(h)) = h \otimes 1, \quad (\varepsilon \otimes id)(\Delta(h)) = 1 \otimes h, \quad (1.2)$$

$$(1 \otimes \Phi)(id \otimes \Delta \otimes id)(\Phi)(\Phi \otimes 1) = (id \otimes id \otimes \Delta)(\Phi)(\Delta \otimes id \otimes id)(\Phi), \quad (1.3)$$

$$(\varepsilon \otimes id \otimes id)(\Phi) = (id \otimes \varepsilon \otimes id)(\Phi) = (id \otimes id \otimes \varepsilon)(\Phi) = 1 \otimes 1 \otimes 1. \quad (1.4)$$

The map Δ is called the coproduct or the comultiplication, ε the counit and Φ the reassociator. We use the version of Sweedler's sigma notation: $\Delta(h) = h_1 \otimes h_2$, for $h \in H$, with the further convention $(\Delta \otimes id)(\Delta(h)) = h_{(1,1)} \otimes h_{(1,2)} \otimes h_2$ and $(id \otimes \Delta)(\Delta(h)) = h_1 \otimes h_{(2,1)} \otimes h_{(2,2)}$. We denote the tensor components of Φ by capital letters and those of Φ^{-1} by small letters:

$$\Phi = X^1 \otimes X^2 \otimes X^3 = T^1 \otimes T^2 \otimes T^3 = Y^1 \otimes Y^2 \otimes Y^3 = \dots$$

$$\Phi^{-1} = x^1 \otimes x^2 \otimes x^3 = t^1 \otimes t^2 \otimes t^3 = y^1 \otimes y^2 \otimes y^3 = \dots$$

The quasi-bialgebra H is called a quasi-Hopf algebra if there exists an anti-automorphism S of the algebra H and elements $\alpha, \beta \in H$ such that, for all $h \in H$, we have:

$$S(h_1)\alpha h_2 = \varepsilon(h)\alpha \quad \text{and} \quad h_1\beta S(h_2) = \varepsilon(h)\beta, \quad (1.5)$$

$$X^1\beta S(X^2)\alpha X^3 = 1 \quad \text{and} \quad S(x^1)\alpha x^2\beta S(x^3) = 1. \quad (1.6)$$

The axioms for a quasi-Hopf algebra imply that $\varepsilon(\alpha)\varepsilon(\beta) = 1$, so, by rescaling α and β , we may assume without loss of generality that $\varepsilon(\alpha) = \varepsilon(\beta) = 1$ and $\varepsilon \circ S = \varepsilon$.

We recall that the definition of a quasi-bialgebra or quasi-Hopf algebra is "twist covariant" in the following sense. An invertible element $F \in H \otimes H$ is called a *gauge transformation* or *twist* if $(\varepsilon \otimes id)(F) = (id \otimes \varepsilon)(F) = 1$. If H is a quasi-bialgebra or a quasi-Hopf algebra and $F = F^1 \otimes F^2 \in H \otimes H$ is a gauge transformation with inverse $F^{-1} = G^1 \otimes G^2$, then we can define a new quasi-bialgebra (respectively quasi-Hopf algebra) H_F by keeping the multiplication, unit, counit (and antipode in the case of a quasi-Hopf algebra) of H and replacing the reassociator, comultiplication and the elements α and β by

$$\Phi_F = (1 \otimes F)(id \otimes \Delta)(F)\Phi(\Delta \otimes id)(F^{-1})(F^{-1} \otimes 1), \quad (1.7)$$

$$\Delta_F(h) = F\Delta(h)F^{-1}, \quad \alpha_F = S(G^1)\alpha G^2, \quad \beta_F = F^1\beta S(F^2). \quad (1.8)$$

The antipode of a Hopf algebra is an anti-coalgebra morphism. For a quasi-Hopf algebra H , we have the following: there exists a gauge transformation $f \in H \otimes H$ such that

$$f\Delta(S(h))f^{-1} = (S \otimes S)(\Delta^{cop}(h)), \quad \text{for all } h \in H. \quad (1.9)$$

The element f can be computed explicitly. First set

$$A^1 \otimes A^2 \otimes A^3 \otimes A^4 = (\Phi \otimes 1)(\Delta \otimes id \otimes id)(\Phi^{-1}), \quad (1.10)$$

$$B^1 \otimes B^2 \otimes B^3 \otimes B^4 = (\Delta \otimes id \otimes id)(\Phi)(\Phi^{-1} \otimes 1), \quad (1.11)$$

and then define $\gamma, \delta \in H \otimes H$ by

$$\gamma = S(A^2)\alpha A^3 \otimes S(A^1)\alpha A^4 \quad \text{and} \quad \delta = B^1\beta S(B^4) \otimes B^2\beta S(B^3). \quad (1.12)$$

Then f and f^{-1} are given by the formulae

$$f = (S \otimes S)(\Delta^{cop}(x^1))\gamma\Delta(x^2\beta S(x^3)), \quad (1.13)$$

$$f^{-1} = \Delta(S(x^1)\alpha x^2)\delta(S \otimes S)(\Delta^{cop}(x^3)). \quad (1.14)$$

Moreover, f satisfies the following relations:

$$f\Delta(\alpha) = \gamma, \quad \Delta(\beta)f^{-1} = \delta, \quad (1.15)$$

and the corresponding twisted reassociator (cf. (1.7)) is given by

$$\Phi_f = S(X^3) \otimes S(X^2) \otimes S(X^1). \quad (1.16)$$

We record also the following relation from [10], for $f^{-1} = g^1 \otimes g^2$ given by (1.14):

$$S^{-1}(\alpha g^2)g^1 = \beta. \quad (1.17)$$

Suppose that $(H, \Delta, \varepsilon, \Phi)$ is a quasi-bialgebra. If U, V, W are left (right) H -modules, define $a_{U,V,W}, \mathbf{a}_{U,V,W} : (U \otimes V) \otimes W \rightarrow U \otimes (V \otimes W)$ by $a_{U,V,W}((u \otimes v) \otimes w) = \Phi \cdot (u \otimes (v \otimes w))$ and $\mathbf{a}_{U,V,W}((u \otimes v) \otimes w) = (u \otimes (v \otimes w)) \cdot \Phi^{-1}$. The category ${}_H\mathcal{M}$ ($\mathcal{M}_H$) of left (right) H -modules becomes a monoidal category (see [19], [21] for terminology) with tensor product $\otimes$ given via Δ , associativity constraints $a_{U,V,W}$ ($\mathbf{a}_{U,V,W}$), unit k as a trivial H -module and the usual left and right unit constraints.

Let again H be a quasi-bialgebra. We say that a k -vector space A is a left H -module algebra if it is an algebra in the monoidal category ${}_H\mathcal{M}$, that is A has a multiplication and a usual unit 1_A satisfying the following conditions:

$$(aa')a'' = (X^1 \cdot a)[(X^2 \cdot a')(X^3 \cdot a'')], \quad (1.18)$$

$$h \cdot (aa') = (h_1 \cdot a)(h_2 \cdot a'), \quad h \cdot 1_A = \varepsilon(h)1_A, \quad (1.19)$$

for all $a, a', a'' \in A$ and $h \in H$, where $h \otimes a \rightarrow h \cdot a$ is the left H -module structure of A . Following [12] we define the smash product $A \# H$ as follows: as vector space $A \# H$ is $A \otimes H$ (elements $a \otimes h$ will be written $a \# h$) with multiplication given by

$$(a \# h)(a' \# h') = (x^1 \cdot a)(x^2 h_1 \cdot a') \# x^3 h_2 h'. \quad (1.20)$$

The smash product $A \# H$ is an associative algebra with unit $1_A \# 1_H$.

If H is a quasi-Hopf algebra, B an associative algebra and $v : H \rightarrow B$ an algebra map, then, following [12], we can introduce on the vector space B a left H -module algebra structure, denoted by B^v in what follows, for which the multiplication, unit and left H -action are:

$$b \star b' = v(X^1)bv(S(x^1 X^2)\alpha x^2 X_1^3)b'v(S(x^3 X_2^3)), \quad \forall b, b' \in B, \quad (1.21)$$

$$1_{B^v} = v(\beta), \quad h \triangleright_v b = v(h_1)bv(S(h_2)), \quad \forall h \in H, b \in B. \quad (1.22)$$

If H is a quasi-Hopf algebra and A is a left H -module algebra, define the following maps:

$$j : H \rightarrow A \# H, \quad j(h) = 1 \# h, \quad \forall h \in H, \quad (1.23)$$

$$i_0 : A \rightarrow A \# H, \quad i_0(a) = x^1 \cdot a \# x^2 \beta S(x^3), \quad \forall a \in A. \quad (1.24)$$

Then, by [12], j is an algebra map and i_0 is a morphism of left H -module algebras from A to $(A \# H)^j$. Moreover, the following universal property of the smash product $A \# H$ holds (see [12], Proposition 2.9): if B is an associative algebra, $v : H \rightarrow B$ is an algebra map and $u : A \rightarrow B^v$ is a morphism of left H -module algebras, then there exists a unique algebra map $u \# v : A \# H \rightarrow B$ such that $(u \# v) \circ i_0 = u$ and $(u \# v) \circ j = v$; this map may be described explicitly as follows:

$$(u \# v)(a \# h) = v(X^1)u(a)v(S(X^2)\alpha X^3h), \quad \forall a \in A, h \in H. \quad (1.25)$$

We record the following relation from [12], which holds in $(A \# H)^j$ for all $a \in A, h \in H$:

$$i_0(a) * j(h) = x^1 \cdot a \# x^2 h S(x^3). \quad (1.26)$$

We recall now the invariance under twisting of the smash product (see for instance [10], [12], [22]). Let H be a quasi-bialgebra, $F \in H \otimes H$ a gauge transformation and A a left H -module algebra. Then we can define a new multiplication on A , by $a \bullet a' = (G^1 \cdot a)(G^2 \cdot a')$, for all $a, a' \in A$, where $F^{-1} = G^1 \otimes G^2$. If we denote by $A_{F^{-1}}$ the resulting structure, then $A_{F^{-1}}$ becomes a left $H_{F^{-1}}$ -module algebra, with the same unit and H -action as for A , and moreover the map $\pi : A \# H \rightarrow A_{F^{-1}} \# H_F, \pi(a \# h) = F^1 \cdot a \# F^2 h$, is an algebra isomorphism.

For further use we need also the notion of right H -module algebra. Let H be a quasi-bialgebra. We say that a k -linear space B is a right H -module algebra if B is an algebra in the monoidal category $\mathcal{M}_H$, i.e. B has a multiplication and a usual unit 1_B satisfying the following conditions:

$$(bb')b'' = (b \cdot x^1)[(b' \cdot x^2)(b'' \cdot x^3)], \quad (1.27)$$

$$(bb') \cdot h = (b \cdot h_1)(b' \cdot h_2), \quad 1_B \cdot h = \varepsilon(h)1_B, \quad (1.28)$$

for all $b, b', b'' \in B$ and $h \in H$, where $b \otimes h \rightarrow b \cdot h$ is the right H -module structure of B .

Recall from [17] the notion of comodule algebra over a quasi-bialgebra.

Definition 1.1 *Let H be a quasi-bialgebra. A unital associative algebra $\mathfrak{A}$ is called a right H -comodule algebra if there exist an algebra morphism $\rho : \mathfrak{A} \rightarrow \mathfrak{A} \otimes H$ and an invertible element $\Phi_\rho \in \mathfrak{A} \otimes H \otimes H$ such that:*

$$\Phi_\rho(\rho \otimes id)(\rho(a)) = (id \otimes \Delta)(\rho(a))\Phi_\rho, \quad \forall a \in \mathfrak{A}, \quad (1.29)$$

$$(1_{\mathfrak{A}} \otimes \Phi)(id \otimes \Delta \otimes id)(\Phi_\rho)(\Phi_\rho \otimes 1_H) = (id \otimes id \otimes \Delta)(\Phi_\rho)(\rho \otimes id \otimes id)(\Phi_\rho), \quad (1.30)$$

$$(id \otimes \varepsilon) \circ \rho = id, \quad (1.31)$$

$$(id \otimes \varepsilon \otimes id)(\Phi_\rho) = (id \otimes id \otimes \varepsilon)(\Phi_\rho) = 1_{\mathfrak{A}} \otimes 1_H. \quad (1.32)$$

Similarly, a unital associative algebra $\mathfrak{B}$ is called a left H -comodule algebra if there exist an algebra morphism $\lambda : \mathfrak{B} \rightarrow H \otimes \mathfrak{B}$ and an invertible element $\Phi_\lambda \in H \otimes H \otimes \mathfrak{B}$ such that:

$$(id \otimes \lambda)(\lambda(b))\Phi_\lambda = \Phi_\lambda(\Delta \otimes id)(\lambda(b)), \quad \forall b \in \mathfrak{B}, \quad (1.33)$$

$$(1_H \otimes \Phi_\lambda)(id \otimes \Delta \otimes id)(\Phi_\lambda)(\Phi_\lambda \otimes 1_{\mathfrak{B}}) = (id \otimes id \otimes \lambda)(\Phi_\lambda)(\Delta \otimes id \otimes id)(\Phi_\lambda), \quad (1.34)$$

$$(\varepsilon \otimes id) \circ \lambda = id, \quad (1.35)$$

$$(id \otimes \varepsilon \otimes id)(\Phi_\lambda) = (\varepsilon \otimes id \otimes id)(\Phi_\lambda) = 1_H \otimes 1_{\mathfrak{B}}. \quad (1.36)$$

When H is a quasi-bialgebra, particular examples of left and right H -comodule algebras are given by $\mathfrak{A} = \mathfrak{B} = H$ and $\rho = \lambda = \Delta$, $\Phi_\rho = \Phi_\lambda = \Phi$. Another basic example of a comodule algebra is provided by the smash product. Namely, if A is a left H -module algebra, then $(A\#H, \rho, \Phi_\rho)$ is a right H -comodule algebra with structures (cf. [12]):

$$\begin{aligned}\rho : A\#H &\rightarrow (A\#H) \otimes H, \quad \rho(a\#h) = (x^1 \cdot a\#x^2h_1) \otimes x^3h_2, \\ \Phi_\rho &= (1\#X^1) \otimes X^2 \otimes X^3 \in (A\#H) \otimes H \otimes H.\end{aligned}$$

For a right H -comodule algebra $(\mathfrak{A}, \rho, \Phi_\rho)$ we will denote, for any $\mathfrak{a} \in \mathfrak{A}$, by $\rho(\mathfrak{a}) = \mathfrak{a}_{<0>} \otimes \mathfrak{a}_{<1>}$, $(\rho \otimes id)(\rho(\mathfrak{a})) = \mathfrak{a}_{<0,0>} \otimes \mathfrak{a}_{<0,1>} \otimes \mathfrak{a}_{<1>}$ etc. Similarly, for a left H -comodule algebra $(\mathfrak{B}, \lambda, \Phi_\lambda)$, if $\mathfrak{b} \in \mathfrak{B}$ then we denote $\lambda(\mathfrak{b}) = \mathfrak{b}_{[-1]} \otimes \mathfrak{b}_{[0]}$, $(id \otimes \lambda)(\lambda(\mathfrak{b})) = \mathfrak{b}_{[-1]} \otimes \mathfrak{b}_{[0,-1]} \otimes \mathfrak{b}_{[0,0]}$ etc. In analogy with the notation for the reassociator Φ of H , we will write

$$\begin{aligned}\Phi_\rho &= \tilde{X}_\rho^1 \otimes \tilde{X}_\rho^2 \otimes \tilde{X}_\rho^3 = \tilde{Y}_\rho^1 \otimes \tilde{Y}_\rho^2 \otimes \tilde{Y}_\rho^3 = \dots \\ \Phi_\rho^{-1} &= \tilde{x}_\rho^1 \otimes \tilde{x}_\rho^2 \otimes \tilde{x}_\rho^3 = \tilde{y}_\rho^1 \otimes \tilde{y}_\rho^2 \otimes \tilde{y}_\rho^3 = \dots\end{aligned}$$

and similarly for the element Φ_λ of a left H -comodule algebra $\mathfrak{B}$.

If $\mathfrak{A}$ is a right H -comodule algebra then we define the elements $\tilde{p}_\rho, \tilde{q}_\rho \in \mathfrak{A} \otimes H$ as follows:

$$\tilde{p}_\rho = \tilde{p}_\rho^1 \otimes \tilde{p}_\rho^2 = \tilde{x}_\rho^1 \otimes \tilde{x}_\rho^2 \beta S(\tilde{x}_\rho^3), \quad \tilde{q}_\rho = \tilde{q}_\rho^1 \otimes \tilde{q}_\rho^2 = \tilde{X}_\rho^1 \otimes S^{-1}(\alpha \tilde{X}_\rho^3) \tilde{X}_\rho^2. \quad (1.37)$$

By [17, Lemma 9.1], we have the following relations, for all $\mathfrak{a} \in \mathfrak{A}$:

$$\rho(\mathfrak{a}_{<0>}) \tilde{p}_\rho[1_{\mathfrak{A}} \otimes S(\mathfrak{a}_{<1>})] = \tilde{p}_\rho[\mathfrak{a} \otimes 1_H], \quad (1.38)$$

$$[1_{\mathfrak{A}} \otimes S^{-1}(\mathfrak{a}_{<1>})] \tilde{q}_\rho \rho(\mathfrak{a}_{<0>}) = [\mathfrak{a} \otimes 1_H] \tilde{q}_\rho, \quad (1.39)$$

$$\rho(\tilde{q}_\rho^1) \tilde{p}_\rho[1_{\mathfrak{A}} \otimes S(\tilde{q}_\rho^2)] = 1_{\mathfrak{A}} \otimes 1_H, \quad (1.40)$$

$$[1_{\mathfrak{A}} \otimes S^{-1}(\tilde{p}_\rho^2)] \tilde{q}_\rho \rho(\tilde{p}_\rho^1) = 1_{\mathfrak{A}} \otimes 1_H, \quad (1.41)$$

$$\begin{aligned}\Phi_\rho(\rho \otimes id_H)(\tilde{p}_\rho)(\tilde{p}_\rho \otimes id_H) \\ = (id_{\mathfrak{A}} \otimes \Delta)(\rho(\tilde{x}_\rho^1) \tilde{p}_\rho)(1_{\mathfrak{A}} \otimes g^1 S(\tilde{x}_\rho^3) \otimes g^2 S(\tilde{x}_\rho^2)),\end{aligned} \quad (1.42)$$

$$\begin{aligned}(\tilde{q}_\rho \otimes 1_H)(\rho \otimes id_H)(\tilde{q}_\rho) \Phi_\rho^{-1} \\ = [1_{\mathfrak{A}} \otimes S^{-1}(f^2 \tilde{X}_\rho^3) \otimes S^{-1}(f^1 \tilde{X}_\rho^2)](id_{\mathfrak{A}} \otimes \Delta)(\tilde{q}_\rho \rho(\tilde{X}_\rho^1)),\end{aligned} \quad (1.43)$$

where $f^{-1} = g^1 \otimes g^2$ is given by (1.14). If $\mathfrak{A}$ is H itself, the elements $\tilde{p}_\rho, \tilde{q}_\rho \in H \otimes H$ are denoted by p_R and q_R and are given by the formulae

$$p_R = p^1 \otimes p^2 = x^1 \otimes x^2 \beta S(x^3), \quad q_R = q^1 \otimes q^2 = X^1 \otimes S^{-1}(\alpha X^3) X^2. \quad (1.44)$$

The next definition appeared in [17] under the name "quasi-commuting pair of H -coactions".

Definition 1.2 *Let H be a quasi-bialgebra. By an H -bicomodule algebra $\mathbb{A}$ we mean a quintuple $(\lambda, \rho, \Phi_\lambda, \Phi_\rho, \Phi_{\lambda,\rho})$, where λ and ρ are left and right H -coactions on $\mathbb{A}$, respectively, and where $\Phi_\lambda \in H \otimes H \otimes \mathbb{A}$, $\Phi_\rho \in \mathbb{A} \otimes H \otimes H$ and $\Phi_{\lambda,\rho} \in H \otimes \mathbb{A} \otimes H$ are invertible elements, such that $(\mathbb{A}, \lambda, \Phi_\lambda)$ is a left H -comodule algebra, $(\mathbb{A}, \rho, \Phi_\rho)$ is a right H -comodule algebra and the following compatibility relations hold:*

$$\Phi_{\lambda,\rho}(\lambda \otimes id)(\rho(u)) = (id \otimes \rho)(\lambda(u)) \Phi_{\lambda,\rho}, \quad \forall u \in \mathbb{A}, \quad (1.45)$$

$$(1_H \otimes \Phi_{\lambda,\rho})(id \otimes \lambda \otimes id)(\Phi_{\lambda,\rho})(\Phi_\lambda \otimes 1_H) = (id \otimes id \otimes \rho)(\Phi_\lambda)(\Delta \otimes id \otimes id)(\Phi_{\lambda,\rho}), \quad (1.46)$$

$$(1_H \otimes \Phi_\rho)(id \otimes \rho \otimes id)(\Phi_{\lambda,\rho})(\Phi_{\lambda,\rho} \otimes 1_H) = (id \otimes id \otimes \Delta)(\Phi_{\lambda,\rho})(\lambda \otimes id \otimes id)(\Phi_\rho). \quad (1.47)$$

As pointed out in [17], if $\mathbb{A}$ is a bicomodule algebra then, in addition, we have that

$$(id_H \otimes id_{\mathbb{A}} \otimes \varepsilon)(\Phi_{\lambda,\rho}) = 1_H \otimes 1_{\mathbb{A}}, \quad (\varepsilon \otimes id_{\mathbb{A}} \otimes id_H)(\Phi_{\lambda,\rho}) = 1_{\mathbb{A}} \otimes 1_H. \quad (1.48)$$

An example of a bicomodule algebra is $\mathbb{A} = H$, $\lambda = \rho = \Delta$ and $\Phi_\lambda = \Phi_\rho = \Phi_{\lambda,\rho} = \Phi$. For the left and right comodule algebra structures of $\mathbb{A}$ we will keep the same notation as above. We also denote $\Phi_{\lambda,\rho} = \Theta^1 \otimes \Theta^2 \otimes \Theta^3 = \bar{\Theta}^1 \otimes \bar{\Theta}^2 \otimes \bar{\Theta}^3$ and $\Phi_{\lambda,\rho}^{-1} = \theta^1 \otimes \theta^2 \otimes \theta^3 = \bar{\theta}^1 \otimes \bar{\theta}^2 \otimes \bar{\theta}^3$.

Let us denote by ${}_H\mathcal{M}_H$ the category of H -bimodules; it is also a monoidal category, the associativity constraints being given by $\mathbf{a}'_{U,V,W} : (U \otimes V) \otimes W \rightarrow U \otimes (V \otimes W)$, $\mathbf{a}'_{U,V,W}((u \otimes v) \otimes w) = \Phi \cdot (u \otimes (v \otimes w)) \cdot \Phi^{-1}$, for $U, V, W \in {}_H\mathcal{M}_H$ and $u \in U$, $v \in V$, $w \in W$. Therefore, we can define algebras in the category of H -bimodules, which will be called H -bimodule algebras. Namely, a k -vector space $\mathcal{A}$ is an H -bimodule algebra if $\mathcal{A}$ is an H -bimodule (denote the actions by $h \cdot \varphi$ and $\varphi \cdot h$, for $h \in H$ and $\varphi \in \mathcal{A}$) with a multiplication and a usual unit $1_{\mathcal{A}}$ such that:

$$(\varphi\psi)\xi = (X^1 \cdot \varphi \cdot x^1)[(X^2 \cdot \psi \cdot x^2)(X^3 \cdot \xi \cdot x^3)], \quad \forall \varphi, \psi, \xi \in \mathcal{A}, \quad (1.49)$$

$$h \cdot (\varphi\psi) = (h_1 \cdot \varphi)(h_2 \cdot \psi), \quad (\varphi\psi) \cdot h = (\varphi \cdot h_1)(\psi \cdot h_2), \quad \forall \varphi, \psi \in \mathcal{A}, h \in H, \quad (1.50)$$

$$h \cdot 1_{\mathcal{A}} = \varepsilon(h)1_{\mathcal{A}}, \quad 1_{\mathcal{A}} \cdot h = \varepsilon(h)1_{\mathcal{A}}, \quad \forall h \in H. \quad (1.51)$$

If H is a quasi-bialgebra then H^* , the linear dual of H , is an H -bimodule with H -actions $\langle h \rightharpoonup \varphi, h' \rangle = \varphi(h'h)$ and $\langle \varphi \leftharpoonup h, h' \rangle = \varphi(hh')$, for all $\varphi \in H^*$, $h, h' \in H$. The convolution $\langle \varphi\psi, h \rangle = \sum \varphi(h_1)\psi(h_2)$, for $\varphi, \psi \in H^*$ and $h \in H$, is a multiplication on H^* , and with this multiplication H^* becomes an H -bimodule algebra. Note that a left (right) H -module algebra becomes also an H -bimodule algebra, with trivial right (left) H -action. Unlike the case of ordinary bialgebras, an H -bimodule algebra is *not* necessarily a left or right H -module algebra, for instance there is no visible left or right H -module algebra structure on H^* .

We recall from [23] the definition of (left) Yetter-Drinfeld modules over a quasi-bialgebra H .

Definition 1.3 A k -linear space M is called a left Yetter-Drinfeld module over H if M is a left H -module (with action denoted by $h \otimes m \mapsto h \cdot m$) and H coacts on M to the left (the coaction is denoted by $\lambda_M : M \rightarrow H \otimes M$, $\lambda_M(m) = m_{(-1)} \otimes m_{(0)}$) such that

$$\begin{aligned} X^1 m_{(-1)} \otimes (X^2 \cdot m_{(0)})_{(-1)} X^3 \otimes (X^2 \cdot m_{(0)})_{(0)} \\ = X^1 (Y^1 \cdot m)_{(-1)_1} Y^2 \otimes X^2 (Y^1 \cdot m)_{(-1)_2} Y^3 \otimes X^3 \cdot (Y^1 \cdot m)_{(0)}, \end{aligned} \quad (1.52)$$

$$\varepsilon(m_{(-1)})m_{(0)} = m, \quad (1.53)$$

$$h_1 m_{(-1)} \otimes h_2 \cdot m_{(0)} = (h_1 \cdot m)_{(-1)} h_2 \otimes (h_1 \cdot m)_{(0)}, \quad (1.54)$$

for all $m \in M$ and $h \in H$. The category ${}^H_H\mathcal{YD}$ consists of such objects, the morphisms in the category being the H -linear maps intertwining the H -coactions.

The category ${}^H_H\mathcal{YD}$ is (pre) braided monoidal; explicitly, if (M, λ_M) and (N, λ_N) are objects in ${}^H_H\mathcal{YD}$, then $(M \otimes N, \lambda_{M \otimes N})$ is also object in ${}^H_H\mathcal{YD}$, where $M \otimes N$ is a left H -module with action $h \cdot (m \otimes n) = h_1 \cdot m \otimes h_2 \cdot n$, and the coaction $\lambda_{M \otimes N}$ is given by

$$\begin{aligned} \lambda_{M \otimes N}(m \otimes n) = X^1 (x^1 Y^1 \cdot m)_{(-1)} x^2 (Y^2 \cdot n)_{(-1)} Y^3 \otimes X^2 \cdot (x^1 Y^1 \cdot m)_{(0)} \\ \otimes X^3 x^3 \cdot (Y^2 \cdot n)_{(0)}. \end{aligned}$$

The associativity constraints are the same as in ${}_H\mathcal{M}$, and the (pre) braiding is given by

$$c_{M,N} : M \otimes N \rightarrow N \otimes M, \quad c_{M,N}(m \otimes n) = m_{(-1)} \cdot n \otimes m_{(0)}.$$

Since ${}^H_H\mathcal{YD}$ is a monoidal category, we can speak about algebras in ${}^H_H\mathcal{YD}$. Namely, if A is an object in ${}^H_H\mathcal{YD}$, then A is an algebra in ${}^H_H\mathcal{YD}$ if and only if A is a left H -module algebra and A is a left quasi-comodule algebra, that is its unit and multiplication intertwine the H -coaction λ_A , namely (for all $a, a' \in A$):

$$\lambda_A(1_A) = 1_H \otimes 1_A, \quad (1.55)$$

$$\begin{aligned} \lambda_A(aa') &= X^1(x^1 Y^1 \cdot a)_{(-1)} x^2 (Y^2 \cdot a')_{(-1)} Y^3 \\ &\quad \otimes [X^2 \cdot (x^1 Y^1 \cdot a)_{(0)}] [X^3 x^3 \cdot (Y^2 \cdot a')_{(0)}]. \end{aligned} \quad (1.56)$$

If H is a quasi-Hopf algebra, we can consider the algebra map $id_H : H \rightarrow H$ and then the left H -module algebra H^{id} , which was denoted by H_0 in [12]; its unit is β , the H -action is $h \triangleright h' = h_1 h' S(h_2)$, and the multiplication is $h * h' = X^1 h S(x^1 X^2) \alpha x^2 X_1^3 h' S(x^3 X_2^3)$. Moreover, it was proved in [11] that H_0 becomes an algebra in ${}^H_H\mathcal{YD}$, with coaction $\lambda_{H_0} : H_0 \rightarrow H \otimes H_0$,

$$\lambda_{H_0}(h) := h_{(-1)} \otimes h_{(0)} = X^1 Y_1^1 h_1 g^1 S(q^2 Y_2^2) Y^3 \otimes X^2 Y_2^1 h_2 g^2 S(X^3 q^1 Y_1^2),$$

where $f^{-1} = g^1 \otimes g^2$ is given by (1.14) and $q_R = q^1 \otimes q^2$ is given by (1.44).

Let H be a quasi-bialgebra, A a left H -module algebra and $\mathfrak{B}$ a left H -comodule algebra. Denote by $A \blacktriangleleft \mathfrak{B}$ the k -vector space $A \otimes \mathfrak{B}$ with multiplication:

$$(a \blacktriangleleft \mathfrak{b})(a' \blacktriangleleft \mathfrak{b}') = (\tilde{x}_\lambda^1 \cdot a)(\tilde{x}_\lambda^2 \mathfrak{b}_{[-1]} \cdot a') \blacktriangleleft \tilde{x}_\lambda^3 \mathfrak{b}_{[0]} \mathfrak{b}', \quad \forall a, a' \in A, \mathfrak{b}, \mathfrak{b}' \in \mathfrak{B}.$$

By [9], $A \blacktriangleleft \mathfrak{B}$ is an associative algebra with unit $1_A \blacktriangleleft 1_{\mathfrak{B}}$, called the generalized smash product of A and $\mathfrak{B}$ (it coincides with the usual smash product if $\mathfrak{B} = H$).

We recall the so-called L-R-smash product, introduced in [28] as a generalization of the cocommutative case from [7], [8]. Let H be a quasi-bialgebra, $\mathcal{A}$ an H -bimodule algebra and $\mathbb{A}$ an H -bicomodule algebra. Define on $\mathcal{A} \otimes \mathbb{A}$ the product

$$(\varphi \natural u)(\varphi' \natural u') = (\tilde{x}_\lambda^1 \cdot \varphi \cdot \theta^3 u'_{<1>} \tilde{x}_\rho^2)(\tilde{x}_\lambda^2 u_{[-1]} \theta^1 \cdot \varphi' \cdot \tilde{x}_\rho^3) \natural \tilde{x}_\lambda^3 u_{[0]} \theta^2 u'_{<0>} \tilde{x}_\rho^1 \quad (1.57)$$

for $\varphi, \varphi' \in \mathcal{A}$ and $u, u' \in \mathbb{A}$, where we write $\varphi \natural u$ in place of $\varphi \otimes u$ to distinguish the new algebraic structure. Then this product defines on $\mathcal{A} \otimes \mathbb{A}$ a structure of associative algebra with unit $1_{\mathcal{A}} \natural 1_{\mathbb{A}}$, denoted by $\mathcal{A} \natural \mathbb{A}$ and called the L-R-smash product. Note that if the right H -action on $\mathcal{A}$ is trivial then $\mathcal{A} \natural \mathbb{A}$ coincides with the generalized smash product $\mathcal{A} \blacktriangleleft \mathbb{A}$.

If H is a quasi-Hopf algebra, $\mathcal{A}$ an H -bimodule algebra and $\mathbb{A}$ an H -bicomodule algebra, there exists another associative algebra structure built on $\mathcal{A} \otimes \mathbb{A}$, which was introduced in [13] under the name generalized diagonal crossed product (for $\mathcal{A} = H^*$ it gives the diagonal crossed product introduced by Hausser and Nill in [17]). It is constructed as follows. Define the element

$$\Omega = (\tilde{X}_\rho^1)_{[-1]_1} \tilde{x}_\lambda^1 \theta^1 \otimes (\tilde{X}_\rho^1)_{[-1]_2} \tilde{x}_\lambda^2 \theta_{[-1]}^2 \otimes (\tilde{X}_\rho^1)_{[0]} \tilde{x}_\lambda^3 \theta_{[0]}^2 \otimes S^{-1}(f^1 \tilde{X}_\rho^2 \theta^3) \otimes S^{-1}(f^2 \tilde{X}_\rho^3) \quad (1.58)$$

in $H^{\otimes 2} \otimes \mathbb{A} \otimes H^{\otimes 2}$, where $\Phi_\rho = \tilde{X}_\rho^1 \otimes \tilde{X}_\rho^2 \otimes \tilde{X}_\rho^3$, $\Phi_\lambda^{-1} = \tilde{x}_\lambda^1 \otimes \tilde{x}_\lambda^2 \otimes \tilde{x}_\lambda^3$, $\Phi_{\lambda, \rho}^{-1} = \theta^1 \otimes \theta^2 \otimes \theta^3$ and $f = f^1 \otimes f^2$ is the twist defined in (1.13). Then define a multiplication on $\mathcal{A} \otimes \mathbb{A}$, by:

$$(\varphi \bowtie u)(\varphi' \bowtie u') = (\Omega^1 \cdot \varphi \cdot \Omega^5)(\Omega^2 u_{<0>[-1]} \cdot \varphi' \cdot S^{-1}(u_{<1>}) \Omega^4) \bowtie \Omega^3 u_{<0>[0]} u', \quad (1.59)$$

for all $\varphi, \varphi' \in \mathcal{A}$ and $u, u' \in \mathbb{A}$, where we denoted $\Omega = \Omega^1 \otimes \dots \otimes \Omega^5$. Then this multiplication defines an associative algebra structure with unit $1_{\mathcal{A}} \bowtie 1_{\mathbb{A}}$, which will be denoted by $\mathcal{A} \bowtie \mathbb{A}$. Note that, as in the case of the L-R-smash product, if the right H -action on $\mathcal{A}$ is trivial then

$\mathcal{A} \bowtie \mathbb{A}$ coincides with the generalized smash product $\mathcal{A} \blacktriangleright \triangleleft \mathbb{A}$.

It was proved in [28] that actually $\mathcal{A} \sharp \mathbb{A}$ and $\mathcal{A} \bowtie \mathbb{A}$ are isomorphic as algebras, a pair of algebra isomorphisms $\nu : \mathcal{A} \bowtie \mathbb{A} \rightarrow \mathcal{A} \sharp \mathbb{A}$ and $\nu^{-1} : \mathcal{A} \sharp \mathbb{A} \rightarrow \mathcal{A} \bowtie \mathbb{A}$ being given by

$$\nu(\varphi \bowtie u) = \Theta^1 \cdot \varphi \cdot S^{-1}(\Theta^3) \tilde{q}_\rho^2 \Theta_{<1>}^2 u_{<1>} \sharp \tilde{q}_\rho^1 \Theta_{<0>}^2 u_{<0>}, \quad (1.60)$$

$$\nu^{-1}(\varphi \sharp u) = \theta^1 \cdot \varphi \cdot S^{-1}(\theta^3 u_{<1>} \tilde{p}_\rho^2) \bowtie \theta^2 u_{<0>} \tilde{p}_\rho^1, \quad (1.61)$$

for all $\varphi \in \mathcal{A}$, $u \in \mathbb{A}$, where $\tilde{p}_\rho = \tilde{p}_\rho^1 \otimes \tilde{p}_\rho^2$ and $\tilde{q}_\rho = \tilde{q}_\rho^1 \otimes \tilde{q}_\rho^2$ are the elements given by (1.37).

We recall now several facts about twisted tensor products of algebras in monoidal categories (see for instance [15], [31]). Let $\mathcal{C}$ be a monoidal category with associativity constraints $a_{U,V,W} : (U \otimes V) \otimes W \rightarrow U \otimes (V \otimes W)$ and unit I . If A is an algebra in $\mathcal{C}$ we denote its multiplication by $\mu_A : A \otimes A \rightarrow A$ and its unit by $\eta_A : I \rightarrow A$. Let (A, μ_A, η_A) and (B, μ_B, η_B) be two algebras in $\mathcal{C}$ and $R : B \otimes A \rightarrow A \otimes B$ a morphism in $\mathcal{C}$. We call R a *twisting map* between A and B if the following conditions hold:

$$R \circ (id_B \otimes \eta_A) = \eta_A \otimes id_B, \quad R \circ (\eta_B \otimes id_A) = id_A \otimes \eta_B, \quad (1.62)$$

$$R \circ (\mu_B \otimes id_A) = (id_A \otimes \mu_B) \circ a_{A,B,B} \circ (R \otimes id_B) \circ a_{B,A,B}^{-1} \circ (id_B \otimes R) \circ a_{B,B,A}, \quad (1.63)$$

$$R \circ (id_B \otimes \mu_A) = (\mu_A \otimes id_B) \circ a_{A,A,B}^{-1} \circ (id_A \otimes R) \circ a_{A,B,A} \circ (R \otimes id_A) \circ a_{B,A,A}^{-1}. \quad (1.64)$$

Given such a twisting map, $A \otimes B$ becomes an algebra in $\mathcal{C}$, with multiplication

$$\begin{aligned} \mu &= (\mu_A \otimes \mu_B) \circ a_{A,A,B \otimes B}^{-1} \circ (id_A \otimes a_{A,B,B}) \circ (id_A \otimes R \otimes id_B) \\ &\quad \circ (id_A \otimes a_{B,A,B}^{-1}) \circ a_{A,B,A \otimes B} \end{aligned} \quad (1.65)$$

and unit $\eta = \eta_A \otimes \eta_B$. This algebra structure on $A \otimes B$ will be denoted by $A \otimes_R B$ and will be called the *twisted tensor product* of A and B afforded by the twisting map R . It has moreover the property that the morphisms $i_A := id_A \otimes \eta_B : A \rightarrow A \otimes_R B$ and $i_B := \eta_A \otimes id_B : B \rightarrow A \otimes_R B$ are morphisms of algebras in $\mathcal{C}$.

If $c_{U,V} : U \otimes V \rightarrow V \otimes U$ is a braiding on $\mathcal{C}$, then for any two algebras A and B in $\mathcal{C}$, the map $R = c_{B,A} : B \otimes A \rightarrow A \otimes B$ is a twisting map, and in this case $A \otimes_R B$ is denoted by $A \underline{\otimes} B$ and is called the *braided tensor product* of A and B (relative to the braiding c).

If $A \otimes_R B$ is a twisted tensor product of algebras in $\mathcal{C}$, it has the following universal property (see [6], [14]): if (X, μ_X, η_X) is an algebra in $\mathcal{C}$ and $u : A \rightarrow X$ and $v : B \rightarrow X$ are morphisms of algebras in $\mathcal{C}$ such that

$$\mu_X \circ (u \otimes v) \circ R = \mu_X \circ (v \otimes u), \quad (1.66)$$

then there exists a unique morphism $w : A \otimes_R B \rightarrow X$ of algebras in $\mathcal{C}$ such that $w \circ i_A = u$ and $w \circ i_B = v$. This morphism w is given explicitly by $w = \mu_X \circ (u \otimes v)$.

2 Twisted tensor products of quasialgebras

Proposition 2.1 *Let H be a quasi-bialgebra, $\mathcal{A}$ an H -bimodule algebra and A an algebra in ${}^H_H\mathcal{YD}$. We regard A as an H -bimodule algebra with trivial right H -action. Define the map*

$$R : A \otimes \mathcal{A} \rightarrow \mathcal{A} \otimes A, \quad R(a \otimes \varphi) = a_{(-1)} \cdot \varphi \otimes a_{(0)}, \quad \forall a \in A, \varphi \in \mathcal{A}. \quad (2.1)$$

Then R is a twisting map between $\mathcal{A}$ and A in the monoidal category ${}_H\mathcal{M}_H$. We will denote by $\mathcal{A} \odot A$ the H -bimodule algebra $\mathcal{A} \otimes_R A$, the twisted tensor product of $\mathcal{A}$ and A in ${}_H\mathcal{M}_H$.

Proof. The fact that R is right H -linear is obvious, and the fact that it is left H -linear follows immediately from (1.54), so indeed R is a morphism in ${}_H\mathcal{M}_H$. The relations (1.62) follow immediately from (1.53) and (1.55). The relation (1.63) reduces to (1.56), while the relation (1.64) reduces to (1.52). $\square$

Remark 2.2 *The explicit structure of $\mathcal{A} \odot A$ is the following: the unit is $1_{\mathcal{A}} \otimes 1_A$, the left H -action is $h \cdot (\varphi \otimes a) = h_1 \cdot \varphi \otimes h_2 \cdot a$, the right H -action is $(\varphi \otimes a) \cdot h = \varphi \cdot h \otimes a$, and the multiplication is (by (1.65)):*

$$\begin{aligned} (\varphi \otimes a)(\varphi' \otimes a') &= (y^1 X^1 \cdot \varphi)(y^2 Y^1(x^1 X^2 \cdot a)_{(-1)} x^2 X_1^3 \cdot \varphi') \\ &\quad \otimes (y_1^3 Y^2 \cdot (x^1 X^2 \cdot a)_{(0)})(y_2^3 Y^3 x^3 X_2^3 \cdot a'). \end{aligned}$$

We will be mainly interested in the following particular case of Proposition 2.1.

Corollary 2.3 *Let H be a quasi-Hopf algebra, A a left H -module algebra and H_0 the algebra in ${}_H^H\mathcal{YD}$ as in the Preliminaries. Define the map $R : H_0 \otimes A \rightarrow A \otimes H_0$,*

$$R(h \otimes a) = h_{(-1)} \cdot a \otimes h_{(0)} = X^1 Y_1^1 h_1 g^1 S(q^2 Y_2^2) Y^3 \cdot a \otimes X^2 Y_2^1 h_2 g^2 S(X^3 q^1 Y_1^2). \quad (2.2)$$

Then R is a twisting map between A and H_0 in the monoidal category ${}_H\mathcal{M}$. We will denote by $A \diamond H_0$ the left H -module algebra $A \otimes_R H_0$, the twisted tensor product of A and H_0 in ${}_H\mathcal{M}$. Its unit is $1_A \otimes \beta$, the H -action is $h \cdot (a \otimes h') = h_1 \cdot a \otimes h_2 \triangleright h'$, and the multiplication is

$$\begin{aligned} (a \otimes h)(a' \otimes h') &= (y^1 X^1 \cdot a)(y^2 Y^1(x^1 X^2 \triangleright h)_{(-1)} x^2 X_1^3 \cdot a') \\ &\quad \otimes (y_1^3 Y^2 \triangleright (x^1 X^2 \triangleright h)_{(0)}) * (y_2^3 Y^3 x^3 X_2^3 \triangleright h'). \end{aligned}$$

More generally, if C is a left H -module algebra and A is an algebra in ${}_H^H\mathcal{YD}$, then $C \odot A$ is a left H -module algebra, which will be denoted by $C \diamond A$.

Since the braiding in ${}_H^H\mathcal{YD}$ is given by $m \otimes n \mapsto m_{(-1)} \cdot n \otimes m_{(0)}$, we obtain:

Corollary 2.4 *Let H be a quasi-Hopf algebra and A an algebra in ${}_H^H\mathcal{YD}$. Then the left H -module algebra $A \diamond H_0$ is actually an algebra in ${}_H^H\mathcal{YD}$ and it coincides with the braided tensor product $A \underline{\otimes} H_0$ in ${}_H^H\mathcal{YD}$.*

As another class of examples of twisted tensor products of quasialgebras, we present the quasialgebra analogue of the so-called *Clifford process*, cf. [3], [32]. Let A be a (not necessarily associative) algebra over k with unit 1, let $q \in k$ be a nonzero fixed element and $\sigma : A \rightarrow A$ an involutive unital algebra automorphism. We denote by $C(k, q)$ the 2-dimensional associative algebra $k[v]/(v^2 = q)$. We define the linear map

$$R : C(k, q) \otimes A \rightarrow A \otimes C(k, q), \quad R(1 \otimes a) = a \otimes 1, \quad R(v \otimes a) = \sigma(a) \otimes v, \quad (2.3)$$

for all $a \in A$. The Clifford process associates to the pair (A, σ) a (not necessarily associative) algebra structure built on $A \otimes C(k, q)$, with multiplication

$$(a \otimes 1 + b \otimes v)(c \otimes 1 + d \otimes v) = (ac + qb\sigma(d)) \otimes 1 + (ad + b\sigma(c)) \otimes v, \quad (2.4)$$

for all $a, b, c, d \in A$. This algebra structure is denoted by $\overline{A}$. Moreover, the linear map

$$\overline{\sigma} : \overline{A} \rightarrow \overline{A}, \quad \overline{\sigma}(a \otimes 1 + b \otimes v) = \sigma(a) \otimes 1 - \sigma(b) \otimes v, \quad \forall a, b \in A,$$

is an involutive unital algebra automorphism for $\overline{A}$. It is clear (cf. [3]) that if A is associative then $\overline{A}$ is also associative, as in this case R is a twisting map and $\overline{A}$ is the twisted tensor product $\overline{A} = A \otimes_R C(k, q)$.

Consider now the initial data for a Clifford process, and assume that A is a left module algebra over a quasi-bialgebra H and σ is H -linear. We regard $C(k, q)$ as a left H -module algebra with trivial H -action. Then one can see that the map R given by (2.3) is a twisting map between A and $C(k, q)$ in the monoidal category ${}_H\mathcal{M}$, and so $\overline{A} = A \otimes_R C(k, q)$ is also a left H -module algebra. It is also clear that the extended involutive automorphism $\overline{\sigma}$ is H -linear too.

Dually, a similar result holds if A is a comodule algebra over a coquasi-bialgebra. In particular, if G is a group and A is a G -graded quasialgebra as in [2] and σ is a graded involutive automorphism of A , then $\overline{A}$ is also a G -graded quasialgebra.

We emphasize an important conceptual difference between the associative and quasiassociative versions of the Clifford process. In the associative case (and assuming $\text{char}(k) \neq 2$), as noted in [2], the algebra A becomes a $\mathbb{Z}_2$ -graded algebra and the twisted tensor product $A \otimes_R C(k, q)$ is actually a braided tensor product, i. e. the twisting map R is obtained from the canonical braiding $c(x \otimes y) = (-1)^{|x||y|} y \otimes x$ of the category of $\mathbb{Z}_2$ -graded vector spaces. In the quasiassociative case, the twisting map R given by (2.3) does *not* seem to come from any braiding.

3 An isomorphism $A \diamond H_0 \simeq (A \# H)^j$ of algebras in ${}_H\mathcal{M}$

Lemma 3.1 *Let H be a quasi-Hopf algebra and A a left H -module algebra. Then for the maps j and i_0 given by (1.23) and (1.24) the following relation holds, for all $h \in H$ and $a \in A$:*

$$j(h) * i_0(a) = i_0(Y^1 X_1^1 h_1 g^1 S(T^2 X_2^2) \alpha T^3 X^3 \cdot a) * j(Y^2 X_2^1 h_2 g^2 S(Y^3 T^1 X_1^2)), \quad (3.1)$$

where $f^{-1} = g^1 \otimes g^2$ is given by (1.14) and $*$ is the multiplication in $(A \# H)^j$.

Proof. We compute:

$$\begin{aligned} j(h) * i_0(a) &= (1 \# h) * (x^1 \cdot a \# x^2 \beta S(x^3)) \\ &= (1 \# X^1)(1 \# h)(1 \# S(y^1 X^2) \alpha y^2 X_1^3) \\ &\quad (x^1 \cdot a \# x^2 \beta S(x^3))(1 \# S(y^3 X_2^3)) \\ &= X_1^1 h_1 S(y^1 X^2)_{1\alpha_1} y_1^2 X_{(1,1)}^3 x^1 \cdot a \\ &\quad \# X_2^1 h_2 S(y^1 X^2)_{2\alpha_2} y_2^2 X_{(1,2)}^3 x^2 \beta S(y^3 X_2^3 x^3) \\ (1.1, 1.15, 1.9) \quad &= X_1^1 h_1 g^1 S(y_2^1 X_2^2) \gamma^1 y_1^2 x^1 X_1^3 \cdot a \\ &\quad \# X_2^1 h_2 g^2 S(y_1^1 X_1^2) \gamma^2 y_2^2 x^2 X_{(2,1)}^3 \beta S(X_{(2,2)}^3) S(y^3 x^3) \\ (1.5, 1.12) \quad &= X_1^1 h_1 g^1 S(T^2 t_2^1 y_2^1 X_2^2) \alpha T^3 t^2 y_1^2 x^1 X^3 \cdot a \\ &\quad \# X_2^1 h_2 g^2 S(T^1 t_1^1 y_1^1 X_1^2) \alpha t^3 y_2^2 x^2 \beta S(y^3 x^3) \\ (1.3, 1.1, 1.5) \quad &= X_1^1 h_1 g^1 S(T^2 X_2^2) \alpha T^3 X^3 \cdot a \# X_2^1 h_2 g^2 S(T^1 X_1^2) S(y^1) \alpha y^2 \beta S(y^3) \\ (1.6) \quad &= X_1^1 h_1 g^1 S(T^2 X_2^2) \alpha T^3 X^3 \cdot a \# X_2^1 h_2 g^2 S(T^1 X_1^2) \\ (1.26) \quad &= i_0(Y^1 X_1^1 h_1 g^1 S(T^2 X_2^2) \alpha T^3 X^3 \cdot a) * j(Y^2 X_2^1 h_2 g^2 S(Y^3 T^1 X_1^2)), \end{aligned}$$

finishing the proof. $\square$

Proposition 3.2 *Let H be a quasi-Hopf algebra and A a left H -module algebra. Then the map $\Pi : A \diamond H_0 \rightarrow (A \# H)^j$, $\Pi(a \otimes h) = x^1 \cdot a \# x^2 h S(x^3)$, for all $a \in A$ and $h \in H$, is an isomorphism of left H -module algebras.*

Proof. Note first that Π is bijective, with inverse given by $\Pi^{-1}(a \# h) = X^1 \cdot a \otimes X^2 h S(X^3)$. For proving that Π is a morphism of left H -module algebras, we will use the universal property of $A \diamond H_0$ as a twisted tensor product of algebras in ${}_H\mathcal{M}$. Namely, we know that $i_0 : A \rightarrow (A \# H)^j$ is a morphism of left H -module algebras, and it is easy to see that $j : H_0 \rightarrow (A \# H)^j$ is also a morphism of left H -module algebras. Moreover, the relation (1.66) reduces in this case exactly to (3.1). We can thus use the universal property of $A \diamond H_0$, which provides a morphism $w : A \diamond H_0 \rightarrow (A \# H)^j$ of left H -module algebras, which is moreover given by $w(a \otimes h) = i_0(a) * j(h)$. Relation (1.26) shows that actually we have $w = \Pi$, finishing the proof. $\square$

As an application of Proposition 3.2, we will prove a certain kind of invariance under Drinfeld twisting for the H -module algebra $A \diamond H_0$. We recall first the following results from [27]:

Lemma 3.3 ([27]) *Let H be a quasi-Hopf algebra, B, C associative algebras, $\eta : B \rightarrow C$, $j : H \rightarrow B$, $v : H \rightarrow C$ algebra maps such that $\eta \circ j = v$. Then the map $\eta : B^j \rightarrow C^v$ is a morphism of left H -module algebras.*

Proposition 3.4 ([27]) *Let H be a quasi-Hopf algebra, $F \in H \otimes H$ a gauge transformation, B an associative algebra, $v : H \rightarrow B$ an algebra map, which will be denoted by v_F when considered as a map from H_F to B . Then the map*

$$\psi : (B^v)_{F^{-1}} \rightarrow B^{v_F}, \quad \psi(b) = v(F^1) b v(S(F^2)), \quad \forall b \in B,$$

is an isomorphism of left H_F -module algebras.

We can state now the desired result.

Proposition 3.5 *Let H be a quasi-Hopf algebra, $F \in H \otimes H$ a gauge transformation and A a left H -module algebra. Then $(A \diamond H_0)_{F^{-1}} \simeq A_{F^{-1}} \diamond (H_F)_0$ as left H_F -module algebras.*

Proof. We denote as before by $j : H \rightarrow A \# H$, $j(h) = 1 \# h$, and by j_F the same map when considered as a map from H_F to $A \# H$ or to $A_{F^{-1}} \# H_F$. Since obviously the algebra isomorphism $\pi : A \# H \simeq A_{F^{-1}} \# H_F$, $\pi(a \# h) = F^1 \cdot a \# F^2 h$, satisfies $\pi \circ j_F = j_F$, we obtain that $(A \# H)^{j_F} \simeq (A_{F^{-1}} \# H_F)^{j_F}$ as left H_F -module algebras, by using Lemma 3.3. On the other hand, by Proposition 3.4, we obtain that $((A \# H)^j)_{F^{-1}} \simeq (A \# H)^{j_F}$ as left H_F -module algebras. Now by Proposition 3.2 we have $A \diamond H_0 \simeq (A \# H)^j$ as left H -module algebras and $A_{F^{-1}} \diamond (H_F)_0 \simeq (A_{F^{-1}} \# H_F)^{j_F}$ as left H_F -module algebras. It is also easy to see that if B and C are isomorphic left H -module algebras then $B_{F^{-1}}$ and $C_{F^{-1}}$ are isomorphic left H_F -module algebras. Using all these facts we finally obtain that $(A \diamond H_0)_{F^{-1}}$ and $A_{F^{-1}} \diamond (H_F)_0$ are isomorphic as left H_F -module algebras. $\square$

4 An isomorphism $A \underline{\otimes} H_0 \simeq (A \# H)^j$ of algebras in ${}^H_H\mathcal{YD}$

We have seen in the previous section that, if H is a quasi-Hopf algebra and A is a left H -module algebra, then the map $\Pi : A \diamond H_0 \rightarrow (A \# H)^j$, $\Pi(a \otimes h) = x^1 \cdot a \# x^2 h S(x^3)$, is an isomorphism of left H -module algebras. If we assume that A is moreover an algebra in ${}^H_H\mathcal{YD}$, in which case $A \diamond H_0$ becomes the braided tensor product $A \underline{\otimes} H_0$, an algebra in ${}^H_H\mathcal{YD}$, then we intend to show that $(A \# H)^j$ becomes also an algebra in ${}^H_H\mathcal{YD}$ in a natural way and Π becomes an isomorphism of algebras in ${}^H_H\mathcal{YD}$. We begin with a result of independent interest.

Proposition 4.1 *Let H be a quasi-bialgebra and A an algebra in ${}^H_H\mathcal{YD}$. Then $(A \# H, \lambda, \Phi_\lambda)$ is a left H -comodule algebra, with structures:*

$$\begin{aligned} \lambda : A \# H &\rightarrow H \otimes (A \# H), \quad \lambda(a \# h) = T^1(t^1 \cdot a)_{(-1)} t^2 h_1 \otimes (T^2 \cdot (t^1 \cdot a)_{(0)} \# T^3 t^3 h_2), \\ \Phi_\lambda &= X^1 \otimes X^2 \otimes (1 \# X^3) \in H \otimes H \otimes (A \# H). \end{aligned}$$

Proof. We first check that λ is an algebra map. It is easy to see that λ is unital, so we only check that it is multiplicative. We compute:

$$\begin{aligned} \lambda((a \# h)(a' \# h')) &= \lambda((z^1 \cdot a)(z^2 h_1 \cdot a') \# z^3 h_2 h') \\ &= T^1[(t_1^1 z^1 \cdot a)(t_2^1 z^2 h_1 \cdot a')]_{(-1)} t^2 z_1^3 h_{(2,1)} h'_1 \\ &\quad \otimes T^2 \cdot [(t_1^1 z^1 \cdot a)(t_2^1 z^2 h_1 \cdot a')]_{(0)} \# T^3 t^3 z_2^3 h_{(2,2)} h'_2 \\ &\stackrel{(1.3)}{=} T^1[(t^1 y^1 \cdot a)(t^2 y_1^2 x^1 h_1 \cdot a')]_{(-1)} t^3 y_2^2 x^2 h_{(2,1)} h'_1 \\ &\quad \otimes T^2 \cdot [(t^1 y^1 \cdot a)(t^2 y_1^2 x^1 h_1 \cdot a')]_{(0)} \# T^3 y^3 x^3 h_{(2,2)} h'_2 \\ &\stackrel{(1.56, 1.1)}{=} T^1 X^1(z^1 y^1 \cdot a)_{(-1)} z^2 (y_1^2 h_{(1,1)} x^1 \cdot a')_{(-1)} y_2^2 h_{(1,2)} x^2 h'_1 \\ &\quad \otimes [T_1^2 X^2 \cdot (z^1 y^1 \cdot a)_{(0)}] [T_2^2 X^3 z^3 \cdot (y_1^2 h_{(1,1)} x^1 \cdot a')_{(0)}] \# T^3 y^3 h_2 x^3 h'_2 \\ &\stackrel{(1.54)}{=} T^1 X^1(z^1 y^1 \cdot a)_{(-1)} z^2 y_1^2 h_{(1,1)} (x^1 \cdot a')_{(-1)} x^2 h'_1 \\ &\quad \otimes [T_1^2 X^2 \cdot (z^1 y^1 \cdot a)_{(0)}] [T_2^2 X^3 z^3 y_2^2 h_{(1,2)} \cdot (x^1 \cdot a')_{(0)}] \# T^3 y^3 h_2 x^3 h'_2 \\ &\stackrel{(1.3)}{=} T^1 X^1(y_1^1 z^1 \cdot a)_{(-1)} y_2^1 z^2 Z^1 h_{(1,1)} (x^1 \cdot a')_{(-1)} x^2 h'_1 \\ &\quad \otimes [T_1^2 X^2 \cdot (y_1^1 z^1 \cdot a)_{(0)}] [T_2^2 X^3 y^2 z_1^3 Z^2 h_{(1,2)} \cdot (x^1 \cdot a')_{(0)}] \\ &\quad \# T^3 y^3 z_2^3 Z^3 h_2 x^3 h'_2 \\ &\stackrel{(1.54)}{=} T^1 X^1 y_1^1 (z^1 \cdot a)_{(-1)} z^2 Z^1 h_{(1,1)} (x^1 \cdot a')_{(-1)} x^2 h'_1 \\ &\quad \otimes [T_1^2 X^2 y_2^1 \cdot (z^1 \cdot a)_{(0)}] [T_2^2 X^3 y^2 z_1^3 Z^2 h_{(1,2)} \cdot (x^1 \cdot a')_{(0)}] \\ &\quad \# T^3 y^3 z_2^3 Z^3 h_2 x^3 h'_2 \\ &\stackrel{(1.1, 1.3)}{=} T^1(z^1 \cdot a)_{(-1)} z^2 h_1 Z^1 (x^1 \cdot a')_{(-1)} x^2 h'_1 \otimes [y^1 T^2 \cdot (z^1 \cdot a)_{(0)}] \\ &\quad [y^2 T_1^3 z_1^3 h_{(2,1)} Z^2 \cdot (x^1 \cdot a')_{(0)}] \# y^3 T_2^3 z_2^3 h_{(2,2)} Z^3 x^3 h'_2 \\ &= [T^1(z^1 \cdot a)_{(-1)} z^2 h_1 \otimes (T^2 \cdot (z^1 \cdot a)_{(0)} \# T^3 z^3 h_2)] \\ &\quad [Z^1 (x^1 \cdot a')_{(-1)} x^2 h'_1 \otimes (Z^2 \cdot (x^1 \cdot a')_{(0)} \# Z^3 x^3 h'_2)] \\ &= \lambda(a \# h) \lambda(a' \# h'), \quad q.e.d. \end{aligned}$$

Now we check (1.33). We compute:

$$\begin{aligned} (id \otimes \lambda)(\lambda(a \# h)) \Phi_\lambda &= [T^1(t^1 \cdot a)_{(-1)} t^2 h_1 \otimes X^1(x^1 T^2 \cdot (t^1 \cdot a)_{(0)})_{(-1)} x^2 T_1^3 t_1^3 h_{(2,1)}] \end{aligned}$$

$$\begin{aligned}
& \otimes X^2 \cdot (x^1 T^2 \cdot (t^1 \cdot a)_{(0)})_{(0)} \# X^3 x^3 T_2^3 t_2^3 h_{(2,2)}] \Phi_\lambda \\
(1.3) \quad & [Y^1 Z^1 x_1^1 (t^1 \cdot a)_{(-1)} t^2 h_1 \otimes X^1 (Y_1^2 Z^2 x_2^1 \cdot (t^1 \cdot a)_{(0)})_{(-1)} Y_2^2 Z^3 x^2 t_1^3 h_{(2,1)} \\
& \otimes X^2 \cdot (Y_1^2 Z^2 x_2^1 \cdot (t^1 \cdot a)_{(0)})_{(0)} \# X^3 Y^3 x^3 t_2^3 h_{(2,2)}] \Phi_\lambda \\
(1.54) \quad & Y^1 Z^1 x_1^1 (t^1 \cdot a)_{(-1)} t^2 h_1 T^1 \otimes X^1 Y_1^2 (Z^2 x_2^1 \cdot (t^1 \cdot a)_{(0)})_{(-1)} Z^3 x^2 t_1^3 h_{(2,1)} T^2 \\
& \otimes X^2 Y_2^2 \cdot (Z^2 x_2^1 \cdot (t^1 \cdot a)_{(0)})_{(0)} \# X^3 Y^3 x^3 t_2^3 h_{(2,2)} T^3 \\
(1.1) \quad & Y^1 Z^1 x_1^1 (t^1 \cdot a)_{(-1)} t^2 T^1 h_{(1,1)} \otimes X^1 Y_1^2 (Z^2 x_2^1 \cdot (t^1 \cdot a)_{(0)})_{(-1)} Z^3 x^2 t_1^3 T^2 h_{(1,2)} \\
& \otimes X^2 Y_2^2 \cdot (Z^2 x_2^1 \cdot (t^1 \cdot a)_{(0)})_{(0)} \# X^3 Y^3 x^3 t_2^3 T^3 h_2 \\
(1.3) \quad & Y^1 Z^1 x_1^1 (T_1^1 z^1 t^1 \cdot a)_{(-1)} T_2^1 z^2 t_1^2 h_{(1,1)} \otimes X^1 Y_1^2 (Z^2 x_2^1 \cdot (T_1^1 z^1 t^1 \cdot a)_{(0)})_{(-1)} \\
& Z^3 x^2 T^2 z^3 t_2^2 h_{(1,2)} \otimes X^2 Y_2^2 \cdot (Z^2 x_2^1 \cdot (T_1^1 z^1 t^1 \cdot a)_{(0)})_{(0)} \# X^3 Y^3 x^3 T^3 t^3 h_2 \\
(1.54) \quad & Y^1 Z^1 (z^1 t^1 \cdot a)_{(-1)} z^2 t_1^2 h_{(1,1)} \otimes X^1 Y_1^2 (Z^2 \cdot (z^1 t^1 \cdot a)_{(0)})_{(-1)} Z^3 z^3 t_2^2 h_{(1,2)} \\
& \otimes X^2 Y_2^2 \cdot (Z^2 \cdot (z^1 t^1 \cdot a)_{(0)})_{(0)} \# X^3 Y^3 t^3 h_2 \\
(1.52) \quad & Y^1 Z^1 (t^1 \cdot a)_{(-1)_1} t_1^2 h_{(1,1)} \otimes X^1 Y_1^2 Z^2 (t^1 \cdot a)_{(-1)_2} t_2^2 h_{(1,2)} \\
& \otimes X^2 Y_2^2 Z^3 \cdot (t^1 \cdot a)_{(0)} \# X^3 Y^3 t^3 h_2 \\
(1.3) \quad & X^1 T_1^1 (t^1 \cdot a)_{(-1)_1} t_1^2 h_{(1,1)} \otimes X^2 T_2^1 (t^1 \cdot a)_{(-1)_2} t_2^2 h_{(1,2)} \otimes X_1^3 T^2 \cdot (t^1 \cdot a)_{(0)} \\
& \# X_2^3 T^3 t^3 h_2 \\
& = \Phi_\lambda(\Delta \otimes id)(\lambda(a \# h)), \quad q.e.d.
\end{aligned}$$

The conditions (1.34), (1.35) and (1.36) are easy to check and are left to the reader. $\square$

Definition 4.2 Let H be a quasi-bialgebra and $(\mathfrak{B}, \lambda, \Phi_\lambda)$, $(\mathfrak{C}, \mu, \Phi_\mu)$ two left H -comodule algebras. A morphism of comodule algebras from $\mathfrak{B}$ to $\mathfrak{C}$ is an algebra map $\nu : \mathfrak{B} \rightarrow \mathfrak{C}$ such that $\Phi_\mu = (id_H \otimes id_H \otimes \nu)(\Phi_\lambda)$ and $\mu \circ \nu = (id_H \otimes \nu) \circ \lambda$.

Remark 4.3 If H is a quasi-bialgebra, A is an algebra in ${}^H_H \mathcal{YD}$ and we consider the left H -comodule algebra $A \# H$ as in Proposition 4.1, then one can easily see that the map $j : H \rightarrow A \# H$, $j(h) = 1 \# h$, is a morphism of left H -comodule algebras.

The next result is a generalization of the fact, proved in [11], Proposition 2.5, that H_0 is an algebra in ${}^H_H \mathcal{YD}$; the proof is similar to the one in [11] and will be omitted.

Proposition 4.4 If H is a quasi-Hopf algebra, $(\mathfrak{B}, \lambda, \Phi_\lambda)$ a left H -comodule algebra and $v : H \rightarrow \mathfrak{B}$ a morphism of H -comodule algebras, then $\mathfrak{B}^v$ becomes an algebra in ${}^H_H \mathcal{YD}$ with coaction

$$\mathfrak{B}^v \rightarrow H \otimes \mathfrak{B}^v, \quad \mathfrak{b} \mapsto X^1 Y_1^1 \mathfrak{b}_{[-1]} g^1 S(q^2 Y_2^2) Y^3 \otimes v(X^2 Y_2^1) \mathfrak{b}_{[0]} v(g^2 S(X^3 q^1 Y_1^2)),$$

where $f^{-1} = g^1 \otimes g^2$ is given by (1.14) and $q_R = q^1 \otimes q^2 = Z^1 \otimes S^{-1}(\alpha Z^3) Z^2$. Moreover, the map $v : H_0 \rightarrow \mathfrak{B}^v$ is a morphism of algebras in ${}^H_H \mathcal{YD}$.

As a consequence of these results, we obtain:

Corollary 4.5 Let H be a quasi-Hopf algebra and A an algebra in ${}^H_H \mathcal{YD}$. Then $(A \# H)^j$ becomes an algebra in ${}^H_H \mathcal{YD}$, with coaction $\lambda_{(A \# H)^j} : (A \# H)^j \rightarrow H \otimes (A \# H)^j$,

$$\begin{aligned}
\lambda_{(A \# H)^j}(a \# h) &= X^1 Y_1^1 T^1 (t^1 \cdot a)_{(-1)} t^2 h_1 g^1 S(q^2 Y_2^2) Y^3 \otimes [X_1^2 Y_{(2,1)}^1 T^2 \cdot (t^1 \cdot a)_{(0)} \\
&\quad \# X_2^2 Y_{(2,2)}^1 T^3 t^3 h_2 g^2 S(X^3 q^1 Y_1^2)],
\end{aligned}$$

and the map $j : H_0 \rightarrow (A \# H)^j$ is a morphism of algebras in ${}^H_H \mathcal{YD}$.

Lemma 4.6 *Let H be a quasi-Hopf algebra and A an algebra in ${}^H_H\mathcal{YD}$. Then i_0 is a morphism of algebras in ${}^H_H\mathcal{YD}$ from A to $(A\#H)^j$.*

Proof. We already know that i_0 is a morphism of left H -module algebras from A to $(A\#H)^j$, so the only thing left to prove is that $\lambda_{(A\#H)^j} \circ i_0 = (id_H \otimes i_0) \circ \lambda_A$. We first record the following relations, whose proofs are easy and left to the reader, which hold for the elements $p_R = p^1 \otimes p^2$ and $q_R = q^1 \otimes q^2$ given by (1.44):

$$T^1 p_1^1 \otimes T^2 p_2^1 \otimes T^3 p^2 = y^1 \otimes y_1^2 p^1 \otimes y_2^2 p^2 S(y^3), \quad (4.1)$$

$$Y^1 Z^1 \otimes q^1 Y_1^2 Z^2 \otimes S(q^2 Y_2^2 Z^3) Y^3 = q_1^1 \otimes q_2^1 \otimes S(q^2). \quad (4.2)$$

Now, by denoting $p_R = P^1 \otimes P^2$ another copy of p_R , we compute:

$$\begin{aligned} (\lambda_{(A\#H)^j} \circ i_0)(a) &= \lambda_{(A\#H)^j}(p^1 \cdot a \# p^2) \\ &= X^1 Y_1^1 T^1 (t^1 p^1 \cdot a)_{(-1)} t^2 p_1^2 g^1 S(q^2 Y_2^2) Y^3 \\ &\quad \otimes [X_1^2 Y_{(2,1)}^1 T^2 \cdot (t^1 p^1 \cdot a)_{(0)} \# X_2^2 Y_{(2,2)}^1 T^3 t^3 p_2^2 g^2 S(X^3 q^1 Y_1^2)] \\ (1.42) \quad &= X^1 Y_1^1 T^1 (Z_{(1,1)}^1 p_1^1 P^1 \cdot a)_{(-1)} Z_{(1,2)}^1 p_2^1 P^2 S(q^2 Y_2^2 Z^3) Y^3 \\ &\quad \otimes [X_1^2 Y_{(2,1)}^1 T^2 \cdot (Z_{(1,1)}^1 p_1^1 P^1 \cdot a)_{(0)} \# X_2^2 Y_{(2,2)}^1 T^3 Z_2^1 p^2 S(X^3 q^1 Y_1^2 Z^2)] \\ (1.54, 1.1) \quad &= X^1 T^1 Y_{(1,1)}^1 Z_{(1,1)}^1 p_1^1 (P^1 \cdot a)_{(-1)} P^2 S(q^2 Y_2^2 Z^3) Y^3 \\ &\quad \otimes [X_1^2 T^2 Y_{(1,2)}^1 Z_{(1,2)}^1 p_2^1 \cdot (P^1 \cdot a)_{(0)} \# X_2^2 T^3 Y_2^1 Z_2^1 p^2 S(X^3 q^1 Y_1^2 Z^2)] \\ (4.2) \quad &= X^1 T^1 q_{(1,1,1)}^1 p_1^1 (P^1 \cdot a)_{(-1)} P^2 S(q^2) \\ &\quad \otimes [X_1^2 T^2 q_{(1,1,2)}^1 p_2^1 \cdot (P^1 \cdot a)_{(0)} \# X_2^2 T^3 q_{(1,2)}^1 p^2 S(q_2^1) S(X^3)] \\ (1.38) \quad &= X^1 T^1 p_1^1 q_1^1 (P^1 \cdot a)_{(-1)} P^2 S(q^2) \\ &\quad \otimes [X_1^2 T^2 p_2^1 q_2^1 \cdot (P^1 \cdot a)_{(0)} \# X_2^2 T^3 p^2 S(X^3)] \\ (1.54, 4.1) \quad &= X^1 y^1 (q_1^1 P^1 \cdot a)_{(-1)} q_2^1 P^2 S(q^2) \\ &\quad \otimes [X_1^2 y_1^2 p^1 \cdot (q_1^1 P^1 \cdot a)_{(0)} \# X_2^2 y_2^2 p^2 S(X^3 y^3)] \\ (1.40) \quad &= a_{(-1)} \otimes (p^1 \cdot a_{(0)} \# p^2) \\ &= ((id_H \otimes i_0) \circ \lambda_A)(a), \end{aligned}$$

finishing the proof. $\square$

Theorem 4.7 *Let H be a quasi-Hopf algebra and A an algebra in ${}^H_H\mathcal{YD}$. Then the map $\Pi : A \otimes H_0 \rightarrow (A\#H)^j$, $\Pi(a \otimes h) = x^1 \cdot a \# x^2 h S(x^3)$, is an isomorphism of algebras in ${}^H_H\mathcal{YD}$.*

Proof. We proved that $j : H_0 \rightarrow (A\#H)^j$ and $i_0 : A \rightarrow (A\#H)^j$ are morphisms of algebras in ${}^H_H\mathcal{YD}$, and together with the commutation relation (3.1) this allows to apply the universal property of the twisted tensor product $A \otimes H_0 = A \otimes_R H_0$ in the category ${}^H_H\mathcal{YD}$, obtaining thus a morphism of algebras in ${}^H_H\mathcal{YD}$ between $A \otimes H_0$ and $(A\#H)^j$, ω say, which has to be given by $\omega(a \otimes h) = i_0(a) * j(h)$, that is ω coincides with the map Π , which finishes the proof. $\square$

5 An algebra isomorphism $B^v \# H \simeq B \otimes H$ and applications

Proposition 5.1 *Let H be a quasi-Hopf algebra, B an associative algebra and $v : H \rightarrow B$ an algebra map. Denote by η the algebra map $\eta : H \rightarrow B \otimes H$, $\eta(h) = v(h_1) \otimes h_2$. Define the map*

$$u : B \rightarrow B \otimes H, \quad u(b) = v(x^1)bv(S(x_2^3 X^3)f^1) \otimes x^2 X^1 \beta S(x_1^3 X^2)f^2, \quad (5.1)$$

where $f = f^1 \otimes f^2$ is the Drinfeld twist given by (1.13). Then u is a morphism of left H -module algebras from B^v to $(B \otimes H)^\eta$.

Proof. The fact that $u(v(\beta)) = \eta(\beta)$ follows immediately from (1.15). We check now that u is a morphism of left H -modules:

$$\begin{aligned} h \triangleright_\eta u(b) &= \eta(h_1)u(b)\eta(S(h_2)) \\ &= v(h_{(1,1)})v(x^1)bv(S(x_2^3 X^3)f^1)v(S(h_2)_1) \otimes h_{(1,2)}x^2 X^1 \beta S(x_1^3 X^2)f^2 S(h_2)_2 \\ &\stackrel{(1.9)}{=} v(h_{(1,1)}x^1)bv(S(h_{(2,2)}x_2^3 X^3)f^1) \otimes h_{(1,2)}x^2 X^1 \beta S(h_{(2,1)}x_1^3 X^2)f^2 \\ &\stackrel{(1.1)}{=} v(x^1 h_1)bv(S(x_2^3 h_{(2,2,2)}X^3)f^1) \otimes x^2 h_{(2,1)}X^1 \beta S(x_1^3 h_{(2,2,1)}X^2)f^2 \\ &\stackrel{(1.1, 1.5)}{=} v(x^1 h_1)bv(S(x_2^3 X^3 h_2)f^1) \otimes x^2 X^1 \beta S(x_1^3 X^2)f^2 \\ &= u(h \triangleright_v b), \quad q.e.d. \end{aligned}$$

Now we check that u is multiplicative (we denote by $F = F^1 \otimes F^2$ another copy of f):

$$\begin{aligned} u(b) * u(b') &= \eta(Y^1)u(b)\eta(S(y^1 Y^2)\alpha y^2 Y_1^3)u(b')\eta(S(y^3 Y_2^3)) \\ &= v(Y_1^1 x^1)bv(S(x_2^3 X^3)f^1 S(y^1 Y^2)_1 \alpha_1 y_1^2 Y_{(1,1)}^3 z^1) b' \\ &\quad v(S(z_2^3 Z^3)F^1 S(y^3 Y_2^3)_1) \otimes Y_2^1 x^2 X^1 \beta S(x_1^3 X^2)f^2 S(y^1 Y^2)_2 \alpha_2 \\ &\quad y_2^2 Y_{(1,2)}^3 z^2 Z^1 \beta S(z_1^3 Z^2)F^2 S(y^3 Y_2^3)_2 \\ &\stackrel{(1.9, 1.15)}{=} v(Y_1^1 x^1)bv(S(T^2 t_2^1 y_2^1 Y_2^2 x_2^3 X^3)\alpha T^3 t^2 y_1^2 Y_{(1,1)}^3 z^1) b' v(S(y_2^3 Y_{(2,2)}^3 z_2^3 Z^3)f^1) \\ &\quad \otimes Y_2^1 x^2 X^1 \beta S(T^1 t_1^1 y_1^1 Y_1^2 x_1^3 X^2)\alpha t^3 y_2^2 Y_{(1,2)}^3 z^2 Z^1 \beta S(y_1^3 Y_{(2,1)}^3 z_1^3 Z^2)f^2 \\ &\stackrel{(1.1, 1.5)}{=} v(Y_1^1 x^1)bv(S(T^2 t_2^1 y_2^1 Y_2^2 x_2^3 X^3)\alpha T^3 t^2 y_1^2 z^1 Y_1^3) b' v(S(y_2^3 z_2^3 Z^3 Y_2^3)f^1) \\ &\quad \otimes Y_2^1 x^2 X^1 \beta S(T^1 t_1^1 y_1^1 Y_1^2 x_1^3 X^2)\alpha t^3 y_2^2 z^2 Z^1 \beta S(y_1^3 z_1^3 Z^2)f^2 \\ &\stackrel{(1.3, 1.1, 1.5)}{=} v(Y_1^1 x^1)bv(S(T^2 t_2^1 Y_2^2 x_2^3 X^3)\alpha T^3 t^2 Y_1^3) b' v(S(y_2^3 Z^3 t^3 Y_2^3)f^1) \\ &\quad \otimes Y_2^1 x^2 X^1 \beta S(y^1 T^1 t_1^1 Y_1^2 x_1^3 X^2)\alpha y^2 Z^1 \beta S(y_1^3 Z^2)f^2 \\ &\stackrel{(1.3, 1.1)}{=} v(x^1 Y^1)bv(S(T^2 x_{(1,1,2)}^3 t_2^1 W_2^2 Y_{(2,2)}^2 X^3)\alpha T^3 x_{(1,2)}^3 t^2 W_1^3 Y_1^3) b' \\ &\quad v(S(y_2^3 Z^3 x_2^3 t^3 W_2^3 Y_2^3)f^1) \otimes x^2 W^1 Y_1^2 X^1 \beta \\ &\quad S(y^1 T^1 x_{(1,1,1)}^3 t_1^1 W_1^2 Y_{(2,1)}^2 X^2)\alpha y^2 Z^1 \beta S(y_1^3 Z^2)f^2 \\ &\stackrel{(1.1, 1.5)}{=} v(x^1 Y^1)bv(S(T^2 t_2^1 W_2^2 Y_{(2,2)}^2 X^3)\alpha T^3 t^2 W_1^3 Y_1^3) b' v(S(y_2^3 Z^3 x_2^3 t^3 W_2^3 Y_2^3)f^1) \\ &\quad \otimes x^2 W^1 Y_1^2 X^1 \beta S(y^1 x_1^3 T^1 t_1^1 W_1^2 Y_{(2,1)}^2 X^2)\alpha y^2 Z^1 \beta S(y_1^3 Z^2)f^2 \\ &\stackrel{(1.1, 1.5)}{=} v(x^1 Y^1)bv(S(T^2 t_2^1 W_2^2 X^3 Y^2)\alpha T^3 t^2 W_1^3 Y_1^3) b' v(S(y_2^3 Z^3 x_2^3 t^3 W_2^3 Y_2^3)f^1) \\ &\quad \otimes x^2 W^1 X^1 \beta S(y^1 x_1^3 T^1 t_1^1 W_1^2 X^2)\alpha y^2 Z^1 \beta S(y_1^3 Z^2)f^2 \\ &\stackrel{(1.3, 1.5)}{=} v(x^1 Y^1)bv(S(T^2 t_2^1 z^2 X_1^3 Y^2)\alpha T^3 t^2 z_1^3 X_{(2,1)}^3 Y_1^3) b' \\ &\quad v(S(y_2^3 Z^3 x_2^3 t^3 z_2^3 X_{(2,2)}^3 Y_2^3)f^1) \end{aligned}$$

$$\begin{aligned}
& \otimes x^2 X^1 \beta S(y^1 x_1^3 T^1 t_1^1 z^1 X^2) \alpha y^2 Z^1 \beta S(y_1^3 Z^2) f^2 \\
(1.3, 1.5) \quad & \underset{=}{=} v(x^1 Y^1) b v(S(T^2 t_2^1 z^2 X_1^3 Y^2) \alpha T^3 t^2 z_1^3 X_{(2,1)}^3 Y_1^3) b' \\
& v(S(x_2^3 t^3 z_2^3 X_{(2,2)}^3 Y_2^3) f^1) \otimes x^2 X^1 \beta S(x_1^3 T^1 t_1^1 z^1 X^2) S(y^1) \alpha y^2 \beta S(y^3) f^2 \\
(1.6, 1.3) \quad & \underset{=}{=} v(x^1 Y^1) b v(S(y_1^2 t^1 X_1^3 Y^2) \alpha y_2^2 t^2 X_{(2,1)}^3 Y_1^3) b' v(S(x_2^3 y^3 t^3 X_{(2,2)}^3 Y_2^3) f^1) \\
& \otimes x^2 X^1 \beta S(x_1^3 y^1 X^2) f^2 \\
(1.1, 1.5) \quad & \underset{=}{=} v(x^1 Y^1) b v(S(t^1 Y^2) \alpha t^2 Y_1^3) b' v(S(x_2^3 X^3 t^3 Y_2^3) f^1) \otimes x^2 X^1 \beta S(x_1^3 X^2) f^2 \\
& \underset{=}{=} u(v(Y^1) b v(S(t^1 Y^2) \alpha t^2 Y_1^3) b' v(S(t^3 Y_2^3))) \\
& \underset{=}{=} u(b * b'),
\end{aligned}$$

finishing the proof. $\square$

Corollary 5.2 *The map $\psi : B^v \# H \rightarrow B \otimes H$, given by*

$$\psi(b \# h) = v(X^1 x_1^1) b v(S(X^2 x_1^2) \alpha X^3 x^2 h_1) \otimes x^3 h_2, \quad (5.2)$$

is an algebra map.

Proof. The universal property of the smash product, applied to the maps η and u from Proposition 5.1, provides an algebra map $B^v \# H \rightarrow B \otimes H$, which turns out to be exactly the map ψ given by (5.2), we leave the details to the reader. $\square$

Proposition 5.3 *If H is a quasi-Hopf algebra, B an algebra, $v : H \rightarrow B$ an algebra map, then:*

- (i) *The map $\theta : B \rightarrow B^v \# H$, $\theta(b) = v(z^1) b v(Z^1 \beta S(z^2 Z^2)) \# z^3 Z^3$, is an algebra map.*
- (ii) *The map $\mu : H \rightarrow B^v \# H$, $\mu(h) = v(z^1 Z^1 \beta S(z^2 h_1 Z^2)) \# z^3 h_2 Z^3$, is an algebra map.*
- (iii) *For all $h \in H$ and $b \in B$, the following relation holds:*

$$\theta(b) \mu(h) = \mu(h) \theta(b) = v(z^1) b v(Z^1 \beta S(z^2 h_1 Z^2)) \# z^3 h_2 Z^3.$$

- (iv) *Consequently, the map*

$$\xi : B \otimes H \rightarrow B^v \# H, \quad \xi(b \otimes h) = v(z^1) b v(Z^1 \beta S(z^2 h_1 Z^2)) \# z^3 h_2 Z^3, \quad (5.3)$$

is an algebra map.

Proof. We only prove (i) and leave the rest to the reader. Obviously $\theta(1) = v(\beta) \# 1$, so we only have to check that θ is multiplicative. We compute:

$$\begin{aligned}
\theta(b) \theta(b') &= [x^1 \triangleright_v (v(z^1) b v(Z^1 \beta S(z^2 Z^2)))] * [x^2 z_1^3 Z_1^3 \triangleright_v (v(t^1) b' v(T^1 \beta S(t^2 T^2)))] \\
&\quad \# x^3 z_2^2 Z_2^3 t^3 T^3 \\
&= v(X^1 x_1^1 z^1) b v(Z^1 \beta S(y^1 X^2 x_2^1 z^2 Z^2) \alpha y^2 X_1^3 x_1^2 z_{(1,1)}^3 Z_{(1,1)}^3 t^1) \\
&\quad b' v(T^1 \beta S(y^3 X_2^3 x_2^2 z_{(1,2)}^3 Z_{(1,2)}^3 t^2 T^2)) \# x^3 z_2^2 Z_2^3 t^3 T^3 \\
(1.3, 1.5) \quad & \underset{=}{=} v(X^1 w_1^1 z^1 y^1) b v(Z^1 \beta S(X^2 w_2^2 z^2 y_1^2 x^1 Z^2) \alpha X^3 w^2 z_1^3 y_{(2,1)}^2 x_1^2 Z_{(1,1)}^3 t^1) \\
&\quad b' v(T^1 \beta S(w^3 z_2^3 y_{(2,2)}^2 x_2^2 Z_{(1,2)}^3 t^2 T^2)) \# y^3 x^3 Z_2^3 t^3 T^3 \\
(1.3, 1.5) \quad & \underset{=}{=} v(y^1) b v(Z^1 \beta S(z^1 y_1^2 x^1 Z^2) \alpha z^2 y_{(2,1)}^2 x_1^2 Z_{(1,1)}^3 t^1)
\end{aligned}$$

$$\begin{aligned}
& b'v(T^1\beta S(z^3y_{(2,2)}^2x_2^2Z_{(1,2)}^3t^2T^2))\#y^3x^3Z_2^3t^3T^3 \\
(1.1, 1.5) \quad & v(y^1)bv(Z^1\beta S(z^1x^1Z^2)\alpha z^2x_1^2Z_{(1,1)}^3t^1)b'v(T^1\beta S(y^2z^3x_2^2Z_{(1,2)}^3t^2T^2)) \\
& \#y^3x^3Z_2^3t^3T^3 \\
(1.3, 1.5) \quad & v(y^1)bv(Y^1Z^1\beta S(z^1Y_1^2Z^2)\alpha z^2Y_{(2,1)}^2Z_1^3t^1)b'v(T^1\beta S(y^2z^3Y_{(2,2)}^2Z_2^3t^2T^2)) \\
& \#y^3Y^3t^3T^3 \\
(1.1, 1.5) \quad & v(y^1)bv(Y^1Z^1\beta S(z^1Z^2)\alpha z^2Z_1^3t^1)b'v(T^1\beta S(y^2Y^2z^3Z_2^3t^2T^2))\#y^3Y^3t^3T^3 \\
(1.3, 1.5) \quad & v(y^1)bv(Y^1Z^1\beta S(Z^2)\alpha Z^3t^1)b'v(T^1\beta S(y^2Y^2t^2T^2))\#y^3Y^3t^3T^3 \\
(1.6) \quad & v(y^1)bb'v(T^1\beta S(y^2T^2))\#y^3T^3 \\
& = \\
& = \theta(bb'),
\end{aligned}$$

finishing the proof. $\square$

Theorem 5.4 *The maps ψ and ξ given by (5.2) and respectively (5.3) are inverse to each other, providing thus an algebra isomorphism $B^v\#H \simeq B \otimes H$.*

Proof. We compute:

$$\begin{aligned}
\xi(\psi(b\#h)) &= v(z^1X^1x_1^1)bv(S(X^2x_2^1)\alpha X^3x^2h_1Z^1\beta S(z^2x_1^3h_{(2,1)}Z^2))\#z^3x_2^3h_{(2,2)}Z^3 \\
(1.1, 1.5) \quad & v(z^1X^1x_1^1)bv(S(X^2x_2^1)\alpha X^3x^2Z^1\beta S(z^2x_1^3Z^2))\#z^3x_2^3Z^3h \\
(1.3, 1.5) \quad & v(z^1X^1Y_{(1,1)}^1t_1^1)bv(S(X^2Y_{(1,2)}^1t_2^1)\alpha X^3Y_2^1t^2\beta S(z^2Y^2t^3))\#z^3Y^3h \\
(1.1, 1.5) \quad & v(X^1t_1^1)bv(S(X^2t_2^1)\alpha X^3t^2\beta S(t^3))\#h \\
(1.3, 1.5) \quad & bv(S(t^1)\alpha t^2\beta S(t^3))\#h \\
(1.6) \quad & b\#h, \\
\psi(\xi(b \otimes h)) &= v(X^1x_1^1z^1)bv(Z^1\beta S(X^2x_2^1z^2h_1Z^2)\alpha X^3x^2z_1^3h_{(2,1)}Z_1^3) \otimes x^3z_2^3h_{(2,2)}Z_2^3 \\
(1.3, 1.5) \quad & bv(Z^1\beta S(t^1h_1Z^2)\alpha t^2h_{(2,1)}Z_1^3) \otimes t^3h_{(2,2)}Z_2^3 \\
(1.1, 1.5) \quad & bv(Z^1\beta S(t^1Z^2)\alpha t^2Z_1^3) \otimes ht^3Z_2^3 \\
(1.3, 1.5) \quad & bv(Z^1\beta S(Z^2)\alpha Z^3) \otimes h \\
(1.6) \quad & b \otimes h,
\end{aligned}$$

finishing the proof. $\square$

We present now some applications of Theorem 5.4. We first record two results of independent interest, whose proofs are straightforward and will be omitted.

Lemma 5.5 *If H is a quasi-Hopf algebra, then $(H \otimes H, \lambda, \Phi_\lambda)$ is a left H -comodule algebra, with structures*

$$\begin{aligned}
\lambda : H \otimes H &\rightarrow H \otimes (H \otimes H), \quad \lambda(h' \otimes h) = \Phi \cdot ((\Delta \otimes id_H)(h' \otimes h)) \cdot \Phi^{-1}, \\
\Phi_\lambda &= (id_H \otimes id_H \otimes \Delta)(\Phi) \in H \otimes H \otimes (H \otimes H).
\end{aligned}$$

Moreover, $\Delta : H \rightarrow H \otimes H$ is a morphism of left H -comodule algebras.

Lemma 5.6 *If H is a quasi-Hopf algebra, $\nu : \mathfrak{B} \rightarrow \mathfrak{C}$ is a morphism of left H -comodule algebras and $v : H \rightarrow \mathfrak{B}$, $w : H \rightarrow \mathfrak{C}$ are morphisms of left H -comodule algebras such that $\nu \circ v = w$, then ν is a morphism of algebras in ${}^H_H\mathcal{YD}$ between $\mathfrak{B}^v$ and $\mathfrak{C}^w$, where $\mathfrak{B}^v$ and $\mathfrak{C}^w$ become algebras in ${}^H_H\mathcal{YD}$ as in Proposition 4.4.*

We consider now the particular case $B = H$ and $v = id_H$ in Theorem 5.4, so we have an algebra isomorphism

$$\Psi : H_0 \# H \rightarrow H \otimes H, \quad \Psi(h' \otimes h) = X^1 x_1^1 h' S(X^2 x_2^1) \alpha X^3 x^2 h_1 \otimes x^3 h_2. \quad (5.4)$$

By a slightly longer but also straightforward computation, one can prove the following result:

Lemma 5.7 *If we consider $H_0 \# H$ a left H -comodule algebra as in Proposition 4.1 and $H \otimes H$ a left H -comodule algebra as in Lemma 5.5, then the map Ψ given by (5.4) is an isomorphism of left H -comodule algebras.*

By using (1.5) and (1.6), one can see that $\Psi \circ j = \Delta$, where $j : H \rightarrow H_0 \# H$, $j(h) = 1_{H_0} \# h$. Thus, as a consequence of the above results and of Theorem 4.7, we finally obtain:

Proposition 5.8 *If H is a quasi-Hopf algebra, then $H_0 \otimes H_0 \simeq (H \otimes H)^\Delta$ as algebras in ${}^H_H\mathcal{YD}$.*

As a second application of Theorem 5.4, we will obtain a duality theorem for quasi-Hopf algebras. We recall from [17] the construction of the two-sided crossed product $H \rtimes H^* \ltimes H$ associated to a finite dimensional quasi-Hopf algebra H , which is an associative algebra structure on $H \otimes H^* \otimes H$ with multiplication

$$(h \otimes \varphi \otimes l)(h' \otimes \varphi' \otimes l') = hh'_1 x^1 \otimes (y^1 \rightharpoonup \varphi \leftharpoonup h'_2 x^2)(y^2 l_1 \rightharpoonup \varphi' \leftharpoonup x^3) \otimes y^3 l_2 l',$$

for $h, h', l, l' \in H$ and $\varphi, \varphi' \in H^*$, where $\rightharpoonup$ and $\leftharpoonup$ are the regular actions of H on H^* . We also recall from [9] the construction of the so-called quasi-smash product, denoted by $H \# H^*$, which is a left H -module algebra structure on $H \otimes H^*$, with multiplication and H -action given by

$$\begin{aligned} (h \otimes \varphi)(h' \otimes \varphi') &= hh'_1 x^1 \otimes (\varphi \leftharpoonup h'_2 x^2)(\varphi' \leftharpoonup x^3), \\ h \cdot (h' \otimes \varphi) &= h' \otimes h \rightharpoonup \varphi, \end{aligned}$$

for all $h, h' \in H$ and $\varphi, \varphi' \in H^*$. By [9], the following identification of algebras holds:

$$H \rtimes H^* \ltimes H \equiv (H \# H^*) \# H.$$

By [27], $H \# H^*$ is isomorphic, as left H -module algebras, to $End(H)^v$, where $End(H)$ is regarded as an associative algebra in the usual way and $v : H \rightarrow End(H)$ is a certain algebra map.

All these facts combined with Theorem 5.4 yield the desired duality theorem:

Theorem 5.9 *If H is a finite dimensional quasi-Hopf algebra, then the two-sided crossed product $H \rtimes H^* \ltimes H$ is isomorphic to $End(H) \otimes H$ as associative algebras.*

6 Some iterated products

Let H be a quasi-bialgebra and A an algebra in ${}^H_H\mathcal{YD}$. By Proposition 4.1, $(A \# H, \lambda, \Phi_\lambda)$ is a left H -comodule algebra, and by the Preliminaries $(A \# H, \rho, \Phi_\rho)$ is a right H -comodule algebra.

Proposition 6.1 $(A\#H, \lambda, \rho, \Phi_\lambda, \Phi_\rho, \Phi_{\lambda,\rho})$ is an H -bicomodule algebra, with $\Phi_{\lambda,\rho} = X^1 \otimes (1\#X^2) \otimes X^3 \in H \otimes (A\#H) \otimes H$.

Proof. We first check the relation (1.45). We compute:

$$\begin{aligned}
& \Phi_{\lambda,\rho}(\lambda \otimes id)(\rho(a\#h)) \\
&= \Phi_{\lambda,\rho}(\lambda \otimes id)((x^1 \cdot a\#x^2h_1) \otimes x^3h_2) \\
&= \Phi_{\lambda,\rho}(T^1(t^1x^1 \cdot a)_{(-1)}t^2x_1^2h_{(1,1)} \otimes (T^2 \cdot (t^1x^1 \cdot a)_{(0)}\#T^3t^3x_2^2h_{(1,2)}) \otimes x^3h_2) \\
(1.3) \quad &= X^1T^1(t_1^1x^1 \cdot a)_{(-1)}t_2^1x^2Y^1h_{(1,1)} \otimes (X_1^2T^2 \cdot (t_1^1x^1 \cdot a)_{(0)}\#X_2^2T^3t^2x_1^3Y^2h_{(1,2)}) \\
&\quad \otimes X^3t^3x_2^3Y^3h_2 \\
(1.1, 1.54) \quad &= X^1T^1t_1^1(x^1 \cdot a)_{(-1)}x^2h_1Y^1 \otimes (X_1^2T^2t_2^1 \cdot (x^1 \cdot a)_{(0)}\#X_2^2T^3t^2x_1^3h_{(2,1)}Y^2) \\
&\quad \otimes X^3t^3x_2^3h_{(2,2)}Y^3 \\
(1.3) \quad &= X^1(x^1 \cdot a)_{(-1)}x^2h_1Y^1 \otimes (y^1X^2 \cdot (x^1 \cdot a)_{(0)}\#y^2X_1^3x_1^3h_{(2,1)}Y^2) \\
&\quad \otimes y^3X_2^3x_2^3h_{(2,2)}Y^3 \\
&= (X^1(x^1 \cdot a)_{(-1)}x^2h_1 \otimes (y^1X^2 \cdot (x^1 \cdot a)_{(0)}\#y^2X_1^3x_1^3h_{(2,1)})) \otimes y^3X_2^3x_2^3h_{(2,2)})\Phi_{\lambda,\rho} \\
&= (id \otimes \rho)(X^1(x^1 \cdot a)_{(-1)}x^2h_1 \otimes (X^2 \cdot (x^1 \cdot a)_{(0)}\#X^3x^3h_2))\Phi_{\lambda,\rho} \\
&= (id \otimes \rho)(\lambda(a\#h))\Phi_{\lambda,\rho}, \quad q.e.d.
\end{aligned}$$

Then, using the fact that $\lambda(1\#h) = h_1 \otimes (1\#h_2)$ and $\rho(1\#h) = (1\#h_1) \otimes h_2$, one can see immediately that the conditions (1.46) and (1.47) reduce to the condition (1.3) for Φ . $\square$

Let now again H be a quasi-bialgebra, A an algebra in ${}^H_H\mathcal{YD}$ and $\mathcal{A}$ an H -bimodule algebra. We can consider the H -bicomodule algebra $A\#H$ as above, and then the L-R-smash product $\mathcal{A} \natural (A\#H)$. On the other hand, we can first consider the H -bimodule algebra $\mathcal{A} \odot A$, and then the L-R-smash product $(\mathcal{A} \odot A) \natural H$.

Proposition 6.2 There is an algebra isomorphism $\Psi : \mathcal{A} \natural (A\#H) \simeq (\mathcal{A} \odot A) \natural H$, given by

$$\Psi(\varphi \natural (a\#h)) = (x^1 \cdot \varphi \otimes x^2 \cdot a) \natural x^3h, \quad \forall \varphi \in \mathcal{A}, a \in A, h \in H.$$

Proof. Obviously Ψ is bijective and unital, so we only have to check that it is multiplicative. We first write down the multiplication rule in $\mathcal{A} \natural (A\#H)$:

$$\begin{aligned}
& [\varphi \natural (a\#h)][\varphi' \natural (a'\#h')] \\
&= (\tilde{x}_\lambda^1 \cdot \varphi \cdot \theta^3(a'\#h')_{<1>\tilde{x}_\rho^2})(\tilde{x}_\lambda^2(a\#h)_{[-1]}\theta^1 \cdot \varphi' \cdot \tilde{x}_\rho^3) \natural \tilde{x}_\lambda^3(a\#h)_{[0]}\theta^2(a'\#h')_{<0>\tilde{x}_\rho^1} \\
&= (x^1 \cdot \varphi \cdot t^3w^3h_2'y^2)(x^2Y^1(z^1 \cdot a)_{(-1)}z^2h_1t^1 \cdot \varphi' \cdot y^3) \\
&\quad \natural (1\#x^3)(Y^2 \cdot (z^1 \cdot a)_{(0)}\#Y^3z^3h_2)(1\#t^2)(w^1 \cdot a'\#w^2h_1')(1\#y^1) \\
&= (x^1 \cdot \varphi \cdot t^3w^3h_2'y^2)(x^2Y^1(z^1 \cdot a)_{(-1)}z^2h_1t^1 \cdot \varphi' \cdot y^3) \\
&\quad \natural (v^1x_1^3Y^2 \cdot (z^1 \cdot a)_{(0)})(v^2x_{(2,1)}^3Y_1^3z_1^3h_{(2,1)}t_1^1w^1 \cdot a')\#v^3x_{(2,2)}^3Y_2^3z_2^3h_{(2,2)}t_2^2w^2h_1'y^1,
\end{aligned}$$

where for the second equality we have used the formulae, presented before, giving the H -bicomodule algebra structure of $A\#H$. Now we compute:

$$\begin{aligned}
& \Psi(\varphi \natural (a\#h))\Psi(\varphi' \natural (a'\#h')) \\
&= [(z^1 \cdot \varphi \otimes z^2 \cdot a) \natural z^3h][(w^1 \cdot \varphi' \otimes w^2 \cdot a') \natural w^3h']
\end{aligned}$$

$$\begin{aligned}
&= (x^1 \cdot (z^1 \cdot \varphi \otimes z^2 \cdot a) \cdot t^3 w_2^3 h_2' y^2) (x^2 z_1^3 h_1 t^1 \cdot (w^1 \cdot \varphi' \otimes w^2 \cdot a') \cdot y^3) \\
&\quad \natural x^3 z_2^3 h_2 t^2 w_1^3 h_1' y^1 \\
&= (x_1^1 z^1 \cdot \varphi \cdot t^3 w_2^3 h_2' y^2 \otimes x_2^1 z^2 \cdot a) (x_1^2 z_{(1,1)}^3 h_{(1,1)} t_1^1 w^1 \cdot \varphi' \cdot y^3 \\
&\quad \otimes x_2^2 z_{(1,2)}^3 h_{(1,2)} t_2^1 w^2 \cdot a') \natural x^3 z_2^3 h_2 t^2 w_1^3 h_1' y^1 \\
&= (u^1 X^1 x_1^1 z^1 \cdot \varphi \cdot t^3 w_2^3 h_2' y^2) (u^2 Y^1 (v^1 X^2 x_2^1 z^2 \cdot a)_{(-1)} \\
&\quad v^2 X_1^3 x_1^2 z_{(1,1)}^3 h_{(1,1)} t_1^1 w^1 \cdot \varphi' \cdot y^3) \otimes (u_1^3 Y^2 \cdot (v^1 X^2 x_2^1 z^2 \cdot a)_{(0)}) \\
&\quad (u_2^3 Y^3 v^3 X_2^3 x_2^2 z_{(1,2)}^3 h_{(1,2)} t_2^1 w^2 \cdot a') \natural x^3 z_2^3 h_2 t^2 w_1^3 h_1' y^1 \\
(1.3, 1.1) \quad &= (u^1 x^1 \cdot \varphi \cdot t^3 w_2^3 h_2' y^2) (u^2 Y^1 (x_{(1,1)}^2 v^1 z^1 \cdot a)_{(-1)} x_{(1,2)}^2 v^2 z_1^2 h_{(1,1)} t_1^1 w^1 \cdot \varphi' \cdot y^3) \\
&\quad \otimes (u_1^3 Y^2 \cdot (x_{(1,1)}^2 v^1 z^1 \cdot a)_{(0)}) (u_2^3 Y^3 x_2^2 v^3 z_2^2 h_{(1,2)} t_2^1 w^2 \cdot a') \natural x^3 z_2^3 h_2 t^2 w_1^3 h_1' y^1 \\
(1.3, 1.54) \quad &= (u^1 x^1 \cdot \varphi \cdot t^3 w_2^3 h_2' y^2) (u^2 Y^1 x_{(1,1)}^2 v_1^1 (z^1 \cdot a)_{(-1)} z^2 X^1 h_{(1,1)} t_1^1 w^1 \cdot \varphi' \cdot y^3) \\
&\quad \otimes (u_1^3 Y^2 x_{(1,2)}^2 v_2^1 \cdot (z^1 \cdot a)_{(0)}) (u_2^3 Y^3 x_2^2 v^2 z_1^3 X^2 h_{(1,2)} t_2^1 w^2 \cdot a') \\
&\quad \natural x^3 v^3 z_2^3 X^3 h_2 t^2 w_1^3 h_1' y^1 \\
(1.1, 1.3) \quad &= (u^1 x^1 \cdot \varphi \cdot t^3 w_2^3 h_2' y^2) (u^2 Y^1 x_{(1,1)}^2 v_1^1 (z^1 \cdot a)_{(-1)} z^2 h_1 t^1 \cdot \varphi' \cdot y^3) \\
&\quad \otimes (u_1^3 Y^2 x_{(1,2)}^2 v_2^1 \cdot (z^1 \cdot a)_{(0)}) (u_2^3 Y^3 x_2^2 v^2 z_1^3 h_{(2,1)} t_1^2 w^1 \cdot a') \\
&\quad \natural x^3 v^3 z_2^3 h_{(2,2)} t_2^2 w^2 h_1' y^1 \\
(1.1, 1.3) \quad &= (u_1^1 x^1 \cdot \varphi \cdot t^3 w_2^3 h_2' y^2) (u_2^1 x^2 Y^1 (z^1 \cdot a)_{(-1)} z^2 h_1 t^1 \cdot \varphi' \cdot y^3) \\
&\quad \otimes (u_2^1 v^1 x_1^3 Y^2 \cdot (z^1 \cdot a)_{(0)}) (u_2^2 v^2 x_{(2,1)}^3 Y_1^3 z_1^3 h_{(2,1)} t_1^2 w^1 \cdot a') \\
&\quad \natural u^3 v^3 x_{(2,2)}^3 Y_2^3 z_2^3 h_{(2,2)} t_2^2 w^2 h_1' y^1 \\
&= \Psi([\varphi \natural (a \# h)][\varphi' \natural (a' \# h')]),
\end{aligned}$$

finishing the proof. $\square$

Corollary 6.3 *If H is a quasi-bialgebra, A an algebra in ${}^H_H\mathcal{YD}$ and C a left H -module algebra, then we have an algebra isomorphism $\Psi : C \blacktriangleright\blacktriangleleft (A \# H) \simeq (C \diamond A) \# H$, given by $\Psi(c \blacktriangleright\blacktriangleleft (a \# h)) = (x^1 \cdot c \otimes x^2 \cdot a) \# x^3 h$, for all $c \in C$, $a \in A$, $h \in H$.*

We recall from [13] that, if H is a quasi-bialgebra, C a left H -module algebra and $\mathbb{A}$ an H -bicomodule algebra, then $C \blacktriangleright\blacktriangleleft \mathbb{A}$ becomes a right H -comodule algebra, with structure defined for all $c \in C$ and $u \in \mathbb{A}$ by:

$$\begin{aligned}
\rho : C \blacktriangleright\blacktriangleleft \mathbb{A} &\rightarrow (C \blacktriangleright\blacktriangleleft \mathbb{A}) \otimes H, \quad \rho(c \blacktriangleright\blacktriangleleft u) = (\theta^1 \cdot c \blacktriangleright\blacktriangleleft \theta^2 u_{<0>}) \otimes \theta^3 u_{<1>}, \\
\Phi_\rho &= (1_C \blacktriangleright\blacktriangleleft \tilde{X}_\rho^1) \otimes \tilde{X}_\rho^2 \otimes \tilde{X}_\rho^3 \in (C \blacktriangleright\blacktriangleleft \mathbb{A}) \otimes H \otimes H.
\end{aligned}$$

Proposition 6.4 *The map Ψ in Corollary 6.3 is an isomorphism of right H -comodule algebras.*

Proof. Note first that $C \blacktriangleright\blacktriangleleft (A \# H)$ and $(C \diamond A) \# H$ are indeed right H -comodule algebras, since C and $C \diamond A$ are left H -module algebras and $A \# H$ and H are H -bicomodule algebras. It is easy to see that Ψ respects the reassociators, so we only prove that it intertwines the coactions:

$$\begin{aligned}
& (\Psi \otimes id_H) \circ \rho_C \blacktriangleright_{(A\#H)}(c \blacktriangleright_{(A\#H)}(a\#h)) \\
&= (\Psi \otimes id_H)((\theta^1 \cdot c \blacktriangleright_{\theta^2(a\#h)_{<0>}}) \otimes \theta^3(a\#h)_{<1>}) \\
&= \Psi(z^1 \cdot c \blacktriangleright_{(1\#z^2)}(x^1 \cdot a\#x^2h_1)) \otimes z^3x^3h_2 \\
&= (y^1z^1 \cdot c \otimes y^2z_1^2x^1 \cdot a) \# y^3z_2^2x^2h_1 \otimes z^3x^3h_2 \\
&\stackrel{(1.3)}{=} (y_1^1x^1 \cdot c \otimes y_2^1x^2 \cdot a) \# y^2x_1^3h_1 \otimes y^3x_2^3h_2 \\
&= \rho_{(C \diamond A) \# H}((x^1 \cdot c \otimes x^2 \cdot a) \# x^3h) \\
&= \rho_{(C \diamond A) \# H} \circ \Psi(c \blacktriangleright_{(A\#H)}(a\#h)),
\end{aligned}$$

finishing the proof. $\square$

For our next results we will need the following slight generalization of Proposition 3.2, whose proof is similar and will be omitted.

Proposition 6.5 *Let H be a quasi-Hopf algebra, A a left H -module algebra, $\mathfrak{B}$ a left H -comodule algebra and $v : H \rightarrow \mathfrak{B}$ a morphism of left H -comodule algebras. Then the map $\tilde{v} : H \rightarrow A \blacktriangleright \mathfrak{B}$, $\tilde{v}(h) = 1_A \blacktriangleright v(h)$, is an algebra map and the map $\Pi : A \diamond \mathfrak{B}^v \rightarrow (A \blacktriangleright \mathfrak{B})^{\tilde{v}}$, $\Pi(a \otimes b) = x^1 \cdot a \blacktriangleright v(x^2)b v(S(x^3))$, is an isomorphism of left H -module algebras.*

Proposition 6.6 *Let H be a quasi-Hopf algebra, A an algebra in ${}^H_H\mathcal{YD}$ and C a left H -module algebra. Then we have an isomorphism of algebras in ${}_H\mathcal{M}$:*

$$(C \diamond A) \diamond H_0 \simeq C \diamond (A \underline{\otimes} H_0), \quad (c \otimes a) \otimes h \mapsto X^1 \cdot c \otimes (X^2 \cdot a \otimes X^3 \triangleright h). \quad (6.1)$$

Proof. We have $(C \diamond A) \diamond H_0 \simeq ((C \diamond A) \# H)^j$ by Proposition 3.2, then $((C \diamond A) \# H)^j \simeq (C \blacktriangleright (A \# H))^j$ by using Corollary 6.3, then $(C \blacktriangleright (A \# H))^j \simeq C \diamond (A \# H)^j$ by Proposition 6.5 and finally $C \diamond (A \# H)^j \simeq C \diamond (A \diamond H_0)$ again by Proposition 3.2. Moreover, one can check that the composition of these four isomorphisms in ${}_H\mathcal{M}$ is exactly the map given by (6.1). $\square$

We recall now some more facts from [13]. Let H be a quasi-bialgebra, A a left H -module algebra and B a right H -module algebra. The two-sided smash product $A \# H \# B$ is the following associative algebra structure on $A \otimes H \otimes B$:

$$(a \# h \# b)(a' \# h' \# b') = (x^1 \cdot a)(x^2 h_1 y^1 \cdot a') \# x^3 h_2 y^2 h'_1 z^1 \# (b \cdot y^3 h'_2 z^2)(b' \cdot z^3),$$

for all $a, a' \in A$, $h, h' \in H$ and $b, b' \in B$. For the particular case when $A = k$, this reduces to the (right-handed) smash product $H \# B$, whose multiplication is

$$(h \# b)(h' \# b') = h h'_1 x^1 \# (b \cdot h'_2 x^2)(b' \cdot x^3),$$

for all $b, b' \in B$, $h, h' \in H$. Moreover, $H \# B$ is a left H -comodule algebra, with structures

$$\begin{aligned}
\lambda : H \# B &\rightarrow H \otimes (H \# B), \quad \lambda(h \# b) = h_1 x^1 \otimes (h_2 x^2 \# b \cdot x^3), \quad \forall h \in H, b \in B, \\
\Phi_\lambda &= X^1 \otimes X^2 \otimes (X^3 \# 1_B) \in H \otimes H \otimes (H \# B),
\end{aligned}$$

and the map $v : H \rightarrow H \# B$, $v(h) = h \# 1$, is a morphism of left H -comodule algebras. Moreover, it was proved in [13] that with respect to this comodule algebra structure of $H \# B$ we have $A \# H \# B = A \blacktriangleright (H \# B)$.

Now, if we assume that H is a quasi-Hopf algebra and if we denote by $j : H \rightarrow A \# H \# B$, $j(h) = 1_A \# h \# 1_B$, which is an algebra map, as a consequence of Proposition 6.5 and of the above facts we obtain the following description of the left H -module algebra $(A \# H \# B)^j$ as a twisted tensor product:

Proposition 6.7 $(A \# H \# B)^j \simeq A \diamond (H \# B)^v$ as left H -module algebras.

We discuss now some different kinds of iterated products, namely in the sense of [18]. We begin by recalling the framework in [18], extended to the case of nontrivial associators. Let $\mathcal{C}$ be a monoidal category, A, B, C algebras in $\mathcal{C}$ and $R_1 : B \otimes A \rightarrow A \otimes B$, $R_2 : C \otimes B \rightarrow B \otimes C$, $R_3 : C \otimes A \rightarrow A \otimes C$ twisting maps in $\mathcal{C}$. Define the maps

$$\begin{aligned} T_1 &: C \otimes (A \otimes B) \rightarrow (A \otimes B) \otimes C, \quad T_2 : (B \otimes C) \otimes A \rightarrow A \otimes (B \otimes C), \\ T_1 &= a_{A,B,C}^{-1} \circ (id_A \otimes R_2) \circ a_{A,C,B} \circ (R_3 \otimes id_B) \circ a_{C,A,B}^{-1}, \\ T_2 &= a_{A,B,C} \circ (R_1 \otimes id_C) \circ a_{B,A,C}^{-1} \circ (id_B \otimes R_3) \circ a_{B,C,A}. \end{aligned}$$

Assume that R_1, R_2, R_3 satisfy the braid equation in $\mathcal{C}$:

$$\begin{aligned} (id_A \otimes R_2) \circ a_{A,C,B} \circ (R_3 \otimes id_B) \circ a_{C,A,B}^{-1} \circ (id_C \otimes R_1) \circ a_{C,B,A} \\ = a_{A,B,C} \circ (R_1 \otimes id_C) \circ a_{B,A,C}^{-1} \circ (id_B \otimes R_3) \circ a_{B,C,A} \circ (R_2 \otimes id_A). \end{aligned} \quad (6.2)$$

Then T_1 is a twisting map between $A \otimes_{R_1} B$ and C , T_2 is a twisting map between A and $B \otimes_{R_2} C$, and moreover $a_{A,B,C}$ is an algebra isomorphism between $(A \otimes_{R_1} B) \otimes_{T_1} C$ and $A \otimes_{T_2} (B \otimes_{R_2} C)$.

We present first a quasialgebra analogue of a result in [18].

Proposition 6.8 *Let H be a quasi-Hopf algebra, A, B two left H -module algebras and C an algebra in ${}^H_H\mathcal{YD}$. Assume that $R_1 : B \otimes A \rightarrow A \otimes B$ is a twisting map in ${}_H\mathcal{M}$ and denote by $R_2 : C \otimes B \rightarrow B \otimes C$ and $R_3 : C \otimes A \rightarrow A \otimes C$ the twisting maps in ${}_H\mathcal{M}$ given by $R_2(c \otimes b) = c_{(-1)} \cdot b \otimes c_{(0)}$ and $R_3(c \otimes a) = c_{(-1)} \cdot a \otimes c_{(0)}$, for all $a \in A, b \in B, c \in C$. Then the twisting maps R_1, R_2, R_3 satisfy the braid equation (6.2) in ${}_H\mathcal{M}$ and the iterated product $(A \otimes_{R_1} B) \otimes_{T_1} C$ coincides with $(A \otimes_{R_1} B) \diamond C$.*

Proof. For $a \in A$ and $b \in B$, we denote $R_1(b \otimes a) = a_{R_1} \otimes b_{R_1}$. We check (6.2):

$$\begin{aligned} & (id_A \otimes R_2) \circ a_{A,C,B} \circ (R_3 \otimes id_B) \circ a_{C,A,B}^{-1} \circ (id_C \otimes R_1) \circ a_{C,B,A}(c \otimes b \otimes a) \\ &= Z^1(x^1 Y^1 \cdot c)_{(-1)} x^2 \cdot (Y^3 \cdot a)_{R_1} \otimes [Z^2 \cdot (x^1 Y^1 \cdot c)_{(0)}]_{(-1)} Z^3 x^3 \cdot (Y^2 \cdot b)_{R_1} \\ & \quad \otimes [Z^2 \cdot (x^1 Y^1 \cdot c)_{(0)}]_{(0)} \\ & \stackrel{(1.52)}{=} X^1(Y^1 \cdot c)_{(-1)_1} \cdot (Y^3 \cdot a)_{R_1} \otimes X^2(Y^1 \cdot c)_{(-1)_2} \cdot (Y^2 \cdot b)_{R_1} \otimes X^3 \cdot (Y^1 \cdot c)_{(0)} \\ &= a_{A,B,C}((Y^1 \cdot c)_{(-1)} \cdot R_1(Y^2 \cdot b \otimes Y^3 \cdot a) \otimes (Y^1 \cdot c)_{(0)}) \\ &= a_{A,B,C}(R_1((Y^1 \cdot c)_{(-1)_1} Y^2 \cdot b \otimes (Y^1 \cdot c)_{(-1)_2} Y^3 \cdot a) \otimes (Y^1 \cdot c)_{(0)}) \\ & \stackrel{(1.52)}{=} a_{A,B,C}(R_1(x^1 X^1 c_{(-1)} \cdot b \otimes x^2 (X^2 \cdot c_{(0)})_{(-1)} X^3 \cdot a) \otimes x^3 \cdot (X^2 \cdot c_{(0)})_{(0)}) \\ &= a_{A,B,C} \circ (R_1 \otimes id_C) \circ a_{B,A,C}^{-1} \circ (id_B \otimes R_3) \circ a_{B,C,A} \circ (R_2 \otimes id_A)(c \otimes b \otimes a). \end{aligned}$$

A similar computation shows that in this case the map T_1 is given by $T_1(c \otimes a \otimes b) = c_{(-1)_1} \cdot a \otimes c_{(-1)_2} \cdot b \otimes c_{(0)}$, which implies immediately that $(A \otimes_{R_1} B) \otimes_{T_1} C = (A \otimes_{R_1} B) \diamond C$. $\square$

Second, we will prove that iterating the quasiassociative version of the Clifford process fits inside the theory developed in [18]. Let A be a left module algebra over a quasi-bialgebra H , let σ be an H -linear involutive unital automorphism of A , let q, s be two nonzero scalars and denote $C(k, q) = k[v]/(v^2 = q)$ and $C(k, s) = k[w]/(w^2 = s)$. Consider the linear maps $R_1 : C(k, q) \otimes A \rightarrow A \otimes C(k, q)$, $R_2 : C(k, s) \otimes C(k, q) \rightarrow C(k, q) \otimes C(k, s)$, $R_3 : C(k, s) \otimes$

$A \rightarrow A \otimes C(k, s)$, given by $R_1(1 \otimes a) = a \otimes 1$, $R_1(v \otimes a) = \sigma(a) \otimes v$, $R_2(1 \otimes 1) = 1 \otimes 1$, $R_2(1 \otimes v) = v \otimes 1$, $R_2(w \otimes 1) = 1 \otimes w$, $R_2(w \otimes v) = -v \otimes w$, $R_3(1 \otimes a) = a \otimes 1$, $R_3(w \otimes a) = \sigma(a) \otimes w$. We consider $C(k, q)$ and $C(k, s)$ as left H -module algebras with trivial H -action, and so R_1, R_2, R_3 become twisting maps in the category ${}_H\mathcal{M}$. One can easily check that they satisfy the braid equation (6.2), so we have the two twisting maps T_1 and T_2 and moreover $(A \otimes_{R_1} C(k, q)) \otimes_{T_1} C(k, s) \equiv A \otimes_{T_2} (C(k, q) \otimes_{R_2} C(k, s))$, because $a_{A, C(k, q), C(k, s)}$ is the identity. One can also check that, if we regard $\bar{A} = A \otimes_{R_1} C(k, q)$ as the Clifford process applied to the pair (A, σ) , then $(A \otimes_{R_1} C(k, q)) \otimes_{T_1} C(k, s)$ is the Clifford process applied to the pair $(\bar{A}, \bar{\sigma})$, where $\bar{\sigma}$ is the extended automorphism given by $\bar{\sigma}(a \otimes 1 + b \otimes v) = \sigma(a) \otimes 1 - \sigma(b) \otimes v$.

Note that in the strictly associative case all these facts were obtained in [3] as an immediate consequence of the fact that the associative Clifford process is given (unlike the quasiassociative one) by a *braided* tensor product of algebras.

7 Universal properties

We have seen in Proposition 3.2 a relation between the algebra $A \# H$ and the quasialgebra $A \diamond H_0$. We analyze now the relation between the universal property of $A \# H$ and the universal property of $A \diamond H_0$ regarded as a twisted tensor product of algebras in ${}_H\mathcal{M}$.

Proposition 7.1 *Let H be a quasi-Hopf algebra, A a left H -module algebra, B an associative algebra, $v : H \rightarrow B$ an algebra map and $u : A \rightarrow B^v$ a morphism of left H -module algebras. Then we have*

$$\mu_{B^v} \circ (u \otimes v) \circ R = \mu_{B^v} \circ (v \otimes u), \quad (7.1)$$

where R is given by (2.2), that is, the input data for the universal property of $A \# H$ is also an input data for the universal property of the twisted tensor product $A \otimes_R H_0$ in the category ${}_H\mathcal{M}$. Moreover, if we denote by $u \# v : A \# H \rightarrow B$ the algebra map provided by the universal property of $A \# H$ and by $w : A \diamond H_0 \rightarrow B^v$ the morphism of left H -module algebras provided by the universal property of $A \diamond H_0 = A \otimes_R H_0$, then we have $(u \# v) \circ \Pi = w$, where Π is the isomorphism from Proposition 3.2.

Proof. We denote by $f^{-1} = g^1 \otimes g^2 = G^1 \otimes G^2$ two copies of the element f^{-1} given by (1.14). We first check (7.1):

$$\begin{aligned} & (\mu_{B^v} \circ (u \otimes v) \circ R)(h \otimes a) \\ &= v(T^1)u(Z^1 X_1^1 h_1 g^1 S(Y^2 X_2^2) \alpha Y^3 X^3 \cdot a) v(S(x^1 T^2) \alpha x^2 T_1^3) \\ & \quad v(Z^2 X_2^1 h_2 g^2 S(Z^3 Y^1 X_1^2)) v(S(x^3 T_2^3)) \\ &= v(T^1 Z_1^1 X_{(1,1)}^1 h_{(1,1)} g_1^1 S(Y^2 X_2^2)_1 \alpha_1 Y_1^3 X_1^3) u(a) \\ & \quad v(S(x^1 T^2 Z_2^1 X_{(1,2)}^1 h_{(1,2)} g_2^1 S(Y^2 X_2^2)_2 \alpha_2 Y_2^3 X_2^3) \\ & \quad \alpha x^2 T_1^3 Z^2 X_2^1 h_2 g^2 S(x^3 T_2^3 Z^3 Y^1 X_1^2)) \\ (1.15, 1.9) \quad &= v(T^1 Z_1^1 X_{(1,1)}^1 h_{(1,1)} g_1^1 G^1 S(W^2 t_2^1 Y_2^2 X_{(2,2)}^2) \alpha W^3 t^2 Y_1^3 X_1^3) u(a) \\ & \quad v(S(x^1 T^2 Z_2^1 X_{(1,2)}^1 h_{(1,2)} g_2^1 G^2 S(W^1 t_1^1 Y_1^2 X_{(2,1)}^2) \alpha t^3 Y_2^3 X_2^3) \\ & \quad \alpha x^2 T_1^3 Z^2 X_2^1 h_2 g^2 S(x^3 T_2^3 Z^3 Y^1 X_1^2)) \\ (1.3, 1.1) \quad &= v(T^1 X_1^1 h_1 Z^1 g_1^1 G^1 S(W^2 t_2^1 Y_2^2 X_{(2,2)}^2) \alpha W^3 t^2 Y_1^3 X_1^3) u(a) \end{aligned}$$

$$\begin{aligned}
& v(S(T_1^2 X_{(2,1)}^1 h_{(2,1)} Z^2 g_1^2 G^2 S(W^1 t_1^1 Y_1^2 X_{(2,1)}^2) \alpha t^3 Y_2^3 X_2^3) \\
& \alpha T_2^2 X_{(2,2)}^1 h_{(2,2)} Z^3 g^2 S(T^3 Y^1 X_1^2)) \\
(1.16) \quad & \underset{=}{=} v(T^1 X_1^1 h_1 G^1 S(W^2 t_2^1 Y_2^2 X_{(2,2)}^2 Z^3) \alpha W^3 t^2 Y_1^3 X_1^3) u(a) \\
& v(S(T_1^2 X_{(2,1)}^1 h_{(2,1)} G_1^2 g^1 S(W^1 t_1^1 Y_1^2 X_{(2,1)}^2 Z^2) \alpha t^3 Y_2^3 X_2^3) \\
& \alpha T_2^2 X_{(2,2)}^1 h_{(2,2)} G_2^2 g^2 S(T^3 Y^1 X_1^2 Z^1)) \\
(1.5) \quad & \underset{=}{=} v(X^1 h S(W^2 t_2^1 Y_2^2 X_{(2,2)}^2 Z^3) \alpha W^3 t^2 Y_1^3 X_1^3) u(a) \\
& v(S(g^1 S(W^1 t_1^1 Y_1^2 X_{(2,1)}^2 Z^2) \alpha t^3 Y_2^3 X_2^3) \alpha g^2 S(Y^1 X_1^2 Z^1)) \\
(1.3, 1.5) \quad & \underset{=}{=} v(X^1 h S(t^1 T^2 Y_2^2 X_{(2,2)}^2 Z^3) \alpha t^2 T_1^3 Y_1^3 X_1^3) u(a) \\
& v(S(g^1 S(T^1 Y_1^2 X_{(2,1)}^2 Z^2) \alpha t^3 T_2^3 Y_2^3 X_2^3) \alpha g^2 S(Y^1 X_1^2 Z^1)) \\
(1.3, 1.1) \quad & \underset{=}{=} v(X^1 h S(t^1 T_1^3 Y^2 X_2^2) \alpha t^2 T_{(2,1)}^3 Y_1^3 X_1^3) u(a) \\
& v(S(g^1 S(T^2 Y_2^1 X_{(1,2)}^2) \alpha t^3 T_{(2,2)}^3 Y_2^3 X_2^3) \alpha g^2 S(T^1 Y_1^1 X_{(1,1)}^2)) \\
(1.1, 1.5) \quad & \underset{=}{=} v(X^1 h S(t^1 Y^2 X_2^2) \alpha t^2 Y_1^3 X_1^3) u(a) \\
& v(S(g^1 S(T^2 Y_2^1 X_{(1,2)}^2) \alpha T^3 t^3 Y_2^3 X_2^3) \alpha g^2 S(T^1 Y_1^1 X_{(1,1)}^2)) \\
(1.9) \quad & \underset{=}{=} v(X^1 h S(t^1 Y^2 X_2^2) \alpha t^2 Y_1^3 X_1^3) u(a) \\
& v(S(S(Y^1 X_1^2)_1 g^1 S(T^2) \alpha T^3 t^3 Y_2^3 X_2^3) \alpha S(Y^1 X_1^2)_2 g^2 S(T^1)) \\
(1.5) \quad & \underset{=}{=} v(X^1 h S(t^1 X^2) \alpha t^2 X_1^3) u(a) v(S(g^1 S(T^2) \alpha T^3 t^3 X_2^3) \alpha g^2 S(T^1)) \\
(1.17, 1.6) \quad & \underset{=}{=} v(X^1 h S(t^1 X^2) \alpha t^2 X_1^3) u(a) v(S(t^3 X_2^3)) \\
& = (\mu_{B^v} \circ (v \otimes u))(h \otimes a).
\end{aligned}$$

Now we check the relation $(u \# v) \circ \Pi = \mu_{B^v} \circ (u \otimes v)$. We compute:

$$\begin{aligned}
((u \# v) \circ \Pi)(a \otimes h) &= (u \# v)(x^1 \cdot a \# x^2 h S(x^3)) \\
(1.25) \quad & \underset{=}{=} v(X^1) u(x^1 \cdot a) v(S(X^2) \alpha X^3 x^2 h S(x^3)) \\
&= v(X^1 x_1^1) u(a) v(S(X^2 x_2^1) \alpha X^3 x^2 h S(x^3)) \\
(1.3, 1.5) \quad & \underset{=}{=} v(X^1) u(a) v(S(x^1 X^2) \alpha x^2 X_1^3 h S(x^3 X_2^3)) \\
&= (\mu_{B^v} \circ (u \otimes v))(a \otimes h),
\end{aligned}$$

finishing the proof. $\square$

Let H be an ordinary Hopf algebra and A a left H -module algebra. We can regard $A \# H$ as a twisted tensor product between the associative algebras A and H , and thus it has a corresponding universal property, which is easily seen to be equivalent to the usual universal property of $A \# H$.

We would like to have a quasi-Hopf analogue of this situation, that is, a universal property of the quasi-Hopf smash product $A \# H$ resembling the universal property of a twisted tensor product. It is not clear *a priori* how to formulate such a universal property, since $A \# H$ itself is not a twisted tensor product, neither of associative algebras nor of quasialgebras.

Our strategy to arrive at such a formulation is the following. We obtain first a universal property of the generalized diagonal crossed product $\mathcal{A} \bowtie \mathbb{A}$, which generalizes (and is inspired by) both the universal property of the diagonal crossed product $H^* \bowtie \mathbb{A}$ over a finite dimensional quasi-Hopf algebra H from [17] and the universal property of a generalized diagonal crossed

product over a Hopf algebra, regarded as a twisted tensor product of associative algebras. Then, since $A\#H$ is a particular case of $\mathcal{A} \bowtie \mathbb{A}$, we will obtain the desired universal property of $A\#H$.

If H is a quasi-Hopf algebra, $\mathcal{A}$ an H -bimodule algebra and $\mathbb{A}$ an H -bicomodule algebra, we denote by $\Gamma : \mathcal{A} \rightarrow \mathcal{A} \bowtie \mathbb{A}$ and $j : \mathbb{A} \rightarrow \mathcal{A} \bowtie \mathbb{A}$ the linear maps given by

$$\Gamma(\varphi) = (\tilde{p}_\rho^1)_{[-1]} \cdot \varphi \cdot S^{-1}(\tilde{p}_\rho^2) \bowtie (\tilde{p}_\rho^1)_{[0]}, \quad j(u) = 1_{\mathcal{A}} \bowtie u,$$

for all $\varphi \in \mathcal{A}$, $u \in \mathbb{A}$, where $\tilde{p}_\rho^1 \otimes \tilde{p}_\rho^2$ is given by (1.37). By [13], the map Γ has the property that $\mathcal{A} \bowtie \mathbb{A}$ is generated as algebra by $\mathbb{A}$ and $\Gamma(\mathcal{A})$, while j is obviously an algebra map.

Proposition 7.2 *Let H be a quasi-Hopf algebra, $\mathcal{A}$ an H -bimodule algebra, $\mathbb{A}$ an H -bicomodule algebra, B an associative algebra, $\gamma : \mathbb{A} \rightarrow B$ an algebra map and $v : \mathcal{A} \rightarrow B$ a linear map such that the following conditions are satisfied:*

$$\gamma(u_{<0>})v(\varphi \cdot u_{<1>}) = v(u_{[-1]} \cdot \varphi)\gamma(u_{[0]}), \quad (7.2)$$

$$v(\varphi\varphi') = \gamma(\tilde{X}_\lambda^1)v(\theta^1 \tilde{X}_\lambda^1 \cdot \varphi \cdot \tilde{X}_\rho^2)\gamma(\theta^2)v(\tilde{X}_\lambda^2 \cdot \varphi' \cdot \tilde{X}_\rho^3\theta^3)\gamma(\tilde{X}_\lambda^3), \quad (7.3)$$

$$v(1_{\mathcal{A}}) = 1_B, \quad (7.4)$$

for all $\varphi, \varphi' \in \mathcal{A}$ and $u \in \mathbb{A}$. Then there exists a unique algebra map $w : \mathcal{A} \bowtie \mathbb{A} \rightarrow B$ such that $w \circ \Gamma = v$ and $w \circ j = \gamma$. Moreover, w is given by the formula

$$w(\varphi \bowtie u) = \gamma(\tilde{q}_\rho^1)v(\varphi \cdot \tilde{q}_\rho^2)\gamma(u), \quad (7.5)$$

for all $\varphi \in \mathcal{A}$ and $u \in \mathbb{A}$, where $\tilde{q}_\rho = \tilde{q}_\rho^1 \otimes \tilde{q}_\rho^2$ is the element given by the formula (1.37).

Proof. We first prove the uniqueness of w . By [13], the map Γ satisfies the relation

$$\varphi \bowtie u = (1_{\mathcal{A}} \bowtie \tilde{q}_\rho^1)\Gamma(\varphi \cdot \tilde{q}_\rho^2)(1_{\mathcal{A}} \bowtie u),$$

for all $\varphi \in \mathcal{A}$ and $u \in \mathbb{A}$, hence we can write

$$\begin{aligned} w(\varphi \bowtie u) &= w(j(\tilde{q}_\rho^1)\Gamma(\varphi \cdot \tilde{q}_\rho^2)j(u)) \\ &= w(j(\tilde{q}_\rho^1))w(\Gamma(\varphi \cdot \tilde{q}_\rho^2))w(j(u)) \\ &= \gamma(\tilde{q}_\rho^1)v(\varphi \cdot \tilde{q}_\rho^2)\gamma(u), \end{aligned}$$

showing that w is indeed unique. We now prove the existence part. Define w by formula (7.5); it is obvious that w is unital and satisfies $w \circ j = \gamma$. We check now that $w \circ \Gamma = v$:

$$\begin{aligned} (w \circ \Gamma)(\varphi) &= w((\tilde{p}_\rho^1)_{[-1]} \cdot \varphi \cdot S^{-1}(\tilde{p}_\rho^2) \bowtie (\tilde{p}_\rho^1)_{[0]}) \\ &= \gamma(\tilde{q}_\rho^1)v((\tilde{p}_\rho^1)_{[-1]} \cdot \varphi \cdot S^{-1}(\tilde{p}_\rho^2)\tilde{q}_\rho^2)\gamma((\tilde{p}_\rho^1)_{[0]}) \\ &\stackrel{(7.2)}{=} \gamma(\tilde{q}_\rho^1)\gamma((\tilde{p}_\rho^1)_{<0>})v(\varphi \cdot S^{-1}(\tilde{p}_\rho^2)\tilde{q}_\rho^2(\tilde{p}_\rho^1)_{<1>}) \\ &\stackrel{(1.41)}{=} v(\varphi). \end{aligned}$$

Thus, the only thing left to prove is that w is multiplicative. We denote by $\tilde{Q}_\rho^1 \otimes \tilde{Q}_\rho^2$ another copy of the element $\tilde{q}_\rho$, and we record the obvious relation

$$\tilde{Q}_\rho^1 \tilde{x}_\rho^1 \otimes S^{-1}(\tilde{x}_\rho^3) \tilde{Q}_\rho^2 \tilde{x}_\rho^2 = 1 \otimes S^{-1}(\alpha). \quad (7.6)$$

Now we compute:

$$\begin{aligned}
& w((\varphi \bowtie u)(\varphi' \bowtie u')) \\
&= \gamma(\tilde{q}_\rho^1)v([\Omega^1 \cdot \varphi \cdot \Omega^5(\tilde{q}_\rho^2)_1][\Omega^2 u_{<0>[-1]} \cdot \varphi' \cdot S^{-1}(u_{<1>})\Omega^4(\tilde{q}_\rho^2)_2])\gamma(\Omega^3 u_{<0>[0]} u') \\
(7.3) \quad &= \gamma(\tilde{q}_\rho^1)\gamma(\tilde{X}_\rho^1)v(\theta^1 \tilde{X}_\lambda^1 \Omega^1 \cdot \varphi \cdot \Omega^5(\tilde{q}_\rho^2)_1 \tilde{X}_\rho^2)\gamma(\theta^2) \\
&= v(\tilde{X}_\lambda^2 \Omega^2 u_{<0>[-1]} \cdot \varphi' \cdot S^{-1}(u_{<1>})\Omega^4(\tilde{q}_\rho^2)_2 \tilde{X}_\rho^3 \theta^3)\gamma(\tilde{X}_\lambda^3)\gamma(\Omega^3 u_{<0>[0]} u') \\
(1.58) \quad &= \gamma(\tilde{q}_\rho^1 \tilde{X}_\rho^1)v(\theta^1 (\tilde{Y}_\rho^1)_{[-1]} \bar{\theta}^1 \cdot \varphi \cdot S^{-1}(f^2 \tilde{Y}_\rho^3)(\tilde{q}_\rho^2)_1 \tilde{X}_\rho^2)\gamma(\theta^2) \\
&= v((\tilde{Y}_\rho^1)_{[0]_{[-1]}} \bar{\theta}_{[-1]}^2 u_{<0>[-1]} \cdot \varphi' \cdot S^{-1}(f^1 \tilde{Y}_\rho^2 \bar{\theta}^3 u_{<1>})(\tilde{q}_\rho^2)_2 \tilde{X}_\rho^3 \theta^3) \\
&= \gamma((\tilde{Y}_\rho^1)_{[0]_{[0]}} \bar{\theta}_{[0]}^2 u_{<0>[0]})\gamma(u') \\
(7.2) \quad &= \gamma(\tilde{q}_\rho^1 \tilde{X}_\rho^1)v(\theta^1 (\tilde{Y}_\rho^1)_{[-1]} \bar{\theta}^1 \cdot \varphi \cdot S^{-1}(f^2 \tilde{Y}_\rho^3)(\tilde{q}_\rho^2)_1 \tilde{X}_\rho^2)\gamma(\theta^2 (\tilde{Y}_\rho^1)_{[0]_{<0>}} \bar{\theta}_{<0>}^2 u_{<0>[<0>]}) \\
&= v(\varphi' \cdot S^{-1}(f^1 \tilde{Y}_\rho^2 \bar{\theta}^3 u_{<1>})(\tilde{q}_\rho^2)_2 \tilde{X}_\rho^3 \theta^3 (\tilde{Y}_\rho^1)_{[0]_{<1>}} \bar{\theta}_{<1>}^2 u_{<0>[<1>]})\gamma(u') \\
(1.45) \quad &= \gamma(\tilde{q}_\rho^1 \tilde{X}_\rho^1)v((\tilde{Y}_\rho^1)_{<0>[-1]} \theta^1 \bar{\theta}^1 \cdot \varphi \cdot S^{-1}(f^2 \tilde{Y}_\rho^3)(\tilde{q}_\rho^2)_1 \tilde{X}_\rho^2)\gamma((\tilde{Y}_\rho^1)_{<0>[0]} \theta^2 \bar{\theta}_{<0>}^2 u_{<0>[<0>]}) \\
&= v(\varphi' \cdot S^{-1}(f^1 \tilde{Y}_\rho^2 \bar{\theta}^3 u_{<1>})(\tilde{q}_\rho^2)_2 \tilde{X}_\rho^3 (\tilde{Y}_\rho^1)_{<1>} \theta^3 \bar{\theta}_{<1>}^2 u_{<0>[<1>]})\gamma(u') \\
(1.47) \quad &= \gamma(\tilde{q}_\rho^1 \tilde{X}_\rho^1)v((\tilde{Y}_\rho^1)_{<0>[-1]} (\tilde{y}_\rho^1)_{[-1]} \theta^1 \cdot \varphi \cdot S^{-1}(f^2 \tilde{Y}_\rho^3)(\tilde{q}_\rho^2)_1 \tilde{X}_\rho^2) \\
&= \gamma((\tilde{Y}_\rho^1)_{<0>[0]} (\tilde{y}_\rho^1)_{[0]} \theta^2 \tilde{Z}_\rho^1 u_{<0>[<0>]}) \\
&= v(\varphi' \cdot S^{-1}(f^1 \tilde{Y}_\rho^2 \tilde{y}_\rho^3 \theta_2^3 \tilde{Z}_\rho^3 u_{<1>})(\tilde{q}_\rho^2)_2 \tilde{X}_\rho^3 (\tilde{Y}_\rho^1)_{<1>} \tilde{y}_\rho^2 \theta_1^3 \tilde{Z}_\rho^2 u_{<0>[<1>]})\gamma(u') \\
(7.2) \quad &= \gamma(\tilde{q}_\rho^1 \tilde{X}_\rho^1 (\tilde{Y}_\rho^1)_{<0>[<0>]} (\tilde{y}_\rho^1)_{<0>} v(\theta^1 \cdot \varphi \cdot S^{-1}(f^2 \tilde{Y}_\rho^3)(\tilde{q}_\rho^2)_1 \tilde{X}_\rho^2 (\tilde{Y}_\rho^1)_{<0>[<1>]} (\tilde{y}_\rho^1)_{<1>})) \\
&= \gamma(\theta^2 \tilde{Z}_\rho^1 u_{<0>[<0>]}) \\
&= v(\varphi' \cdot S^{-1}(f^1 \tilde{Y}_\rho^2 \tilde{y}_\rho^3 \theta_2^3 \tilde{Z}_\rho^3 u_{<1>})(\tilde{q}_\rho^2)_2 \tilde{X}_\rho^3 (\tilde{Y}_\rho^1)_{<1>} \tilde{y}_\rho^2 \theta_1^3 \tilde{Z}_\rho^2 u_{<0>[<1>]})\gamma(u') \\
(1.29, 1.30) \quad &= \gamma(\tilde{q}_\rho^1 (\tilde{Y}_\rho^1)_{<0>} \tilde{y}_\rho^1 \tilde{X}_\rho^1)v(\theta^1 \cdot \varphi \cdot S^{-1}(f^2 \tilde{Y}_\rho^3)(\tilde{q}_\rho^2)_1 (\tilde{Y}_\rho^1)_{<1>} (\tilde{y}_\rho^2)_1 x^1 \tilde{X}_\rho^2) \\
&= \gamma(\theta^2 \tilde{Z}_\rho^1 u_{<0>[<0>]})v(\varphi' \cdot S^{-1}(f^1 \tilde{Y}_\rho^2 \tilde{y}_\rho^3 x^3 (\tilde{X}_\rho^3)_2 \theta_2^3 \tilde{Z}_\rho^3 u_{<1>})) \\
&= (\tilde{q}_\rho^2)_2 (\tilde{Y}_\rho^1)_{<1>} (\tilde{y}_\rho^2)_2 x^2 (\tilde{X}_\rho^3)_1 \theta_1^3 \tilde{Z}_\rho^2 u_{<0>[<1>]})\gamma(u') \\
(1.29, 1.43) \quad &= \gamma(\tilde{q}_\rho^1 (\tilde{Q}_\rho^1)_{<0>} \tilde{x}_\rho^1 \tilde{y}_\rho^1 \tilde{X}_\rho^1)v(\theta^1 \cdot \varphi \cdot \tilde{q}_\rho^2 (\tilde{Q}_\rho^1)_{<1>} \tilde{x}_\rho^2 (\tilde{y}_\rho^2)_1 x^1 \tilde{X}_\rho^2)\gamma(\theta^2 u_{<0>} \tilde{Z}_\rho^1) \\
&= v(\varphi' \cdot S^{-1}(\tilde{y}_\rho^3 x^3 (\tilde{X}_\rho^3)_2 \theta_2^3 u_{<1>} \tilde{Z}_\rho^3) \tilde{Q}_\rho^2 \tilde{x}_\rho^3 (\tilde{y}_\rho^2)_2 x^2 (\tilde{X}_\rho^3)_1 \theta_1^3 u_{<1>} \tilde{Z}_\rho^2)\gamma(u') \\
(1.30) \quad &= \gamma(\tilde{q}_\rho^1 (\tilde{Q}_\rho^1)_{<0>} (\tilde{x}_\rho^1)_{<0>} v(\theta^1 \cdot \varphi \cdot \tilde{q}_\rho^2 (\tilde{Q}_\rho^1)_{<1>} (\tilde{x}_\rho^1)_{<1>}))\gamma(\theta^2 u_{<0>} \tilde{Z}_\rho^1) \\
&= v(\varphi' \cdot S^{-1}(\tilde{x}_\rho^3 \theta_2^3 u_{<1>} \tilde{Z}_\rho^3) \tilde{Q}_\rho^2 \tilde{x}_\rho^2 \theta_1^3 u_{<1>} \tilde{Z}_\rho^2)\gamma(u') \\
(7.6, 1.5) \quad &= \gamma(\tilde{q}_\rho^1)v(\varphi \cdot \tilde{q}_\rho^2)\gamma(u \tilde{Z}_\rho^1)v(\varphi' \cdot S^{-1}(\tilde{Z}_\rho^3)S^{-1}(\alpha) \tilde{Z}_\rho^2)\gamma(u') \\
&= \gamma(\tilde{q}_\rho^1)v(\varphi \cdot \tilde{q}_\rho^2)\gamma(u)\gamma(\tilde{Q}_\rho^1)v(\varphi' \cdot \tilde{Q}_\rho^2)\gamma(u') \\
&= w(\varphi \bowtie u)w(\varphi' \bowtie u'),
\end{aligned}$$

finishing the proof. $\square$

Remark 7.3 This universal property of $\mathcal{A} \bowtie \mathbb{A}$ looks quite different from the universal property of $H^* \bowtie \mathbb{A}$ given in [17], Theorem II, which is formulated in terms of so-called *normal coherent intertwiners*, but for finite dimensional H and $\mathcal{A} = H^*$ the two universal properties are actually equivalent. Indeed, if $T = T^1 \otimes T^2 \in H \otimes B$ is a normal coherent intertwiner as in [17], then we can define the map $v : H^* \rightarrow B$, $v(\varphi) = \varphi(T^1)T^2$, and the three conditions (7.2), (7.3) and (7.4) for v follow from the three conditions in [17] defining a normal coherent intertwiner.

We intend now to prove that the algebra isomorphism $\nu : \mathcal{A} \bowtie \mathbb{A} \rightarrow \mathcal{A} \natural \mathbb{A}$ given by (1.60) may be naturally reobtained by using Proposition 7.2. We define the map

$$\Lambda : \mathcal{A} \rightarrow \mathcal{A} \natural \mathbb{A}, \quad \Lambda(\varphi) = \theta^1 \cdot \varphi \cdot \theta^3 \natural \theta^2. \quad (7.7)$$

We claim that the conditions in Proposition 7.2 are satisfied for $B = \mathcal{A} \natural \mathbb{A}$, $\gamma = j$ and $v = \Lambda$. Indeed, condition (7.2) follows easily from (1.45) and condition (7.4) is trivial, so we only have to check (7.3). We denote by $\tilde{\theta}^1 \otimes \tilde{\theta}^2 \otimes \tilde{\theta}^3$ one more copy of $\Phi_{\lambda, \rho}^{-1}$ and we compute:

$$\begin{aligned} & j(\tilde{X}_\rho^1) \Lambda(\theta^1 \tilde{X}_\lambda^1 \cdot \varphi \cdot \tilde{X}_\rho^2) j(\theta^2) \Lambda(\tilde{X}_\lambda^2 \cdot \varphi' \cdot \tilde{X}_\rho^3 \theta^3) j(\tilde{X}_\lambda^3) \\ &= ((\tilde{X}_\rho^1)_{[-1]} \bar{\theta}^1 \theta^1 \tilde{X}_\lambda^1 \cdot \varphi \cdot \tilde{X}_\rho^2 \bar{\theta}^3 \theta_{<1>}^2 \natural (\tilde{X}_\rho^1)_{[0]} \bar{\theta}^2 \theta_{<0>}^2) \\ & \quad (\tilde{\theta}^1 \tilde{X}_\lambda^2 \cdot \varphi' \cdot \tilde{X}_\rho^3 \theta^3 \bar{\theta}^3 (\tilde{X}_\lambda^3)_{<1>} \natural \bar{\theta}^2 (\tilde{X}_\lambda^3)_{<0>}) \\ (1.47) \quad &= (\theta^1 \tilde{X}_\lambda^1 \cdot \varphi \cdot \theta_1^3 \tilde{Y}_\rho^2 \natural \theta^2 \tilde{Y}_\rho^1) (\bar{\theta}^1 \tilde{X}_\lambda^2 \cdot \varphi' \cdot \theta_2^3 \tilde{Y}_\rho^3 \bar{\theta}^3 (\tilde{X}_\lambda^3)_{<1>} \natural \bar{\theta}^2 (\tilde{X}_\lambda^3)_{<0>}) \\ &= (\tilde{x}_\lambda^1 \theta^1 \tilde{X}_\lambda^1 \cdot \varphi \cdot \theta_1^3 \tilde{Y}_\rho^2 \bar{\theta}^3 \bar{\theta}_{<1>}^2 (\tilde{X}_\lambda^3)_{<0> <1>} \tilde{x}_\rho^2) \\ & \quad (\tilde{x}_\lambda^2 \theta_{[-1]}^2 (\tilde{Y}_\rho^1)_{[-1]} \bar{\theta}^1 \bar{\theta}^1 \tilde{X}_\lambda^2 \cdot \varphi' \cdot \theta_2^3 \tilde{Y}_\rho^3 \bar{\theta}^3 (\tilde{X}_\lambda^3)_{<1>} \tilde{x}_\rho^3 \natural \tilde{x}_\lambda^3 \theta_{[0]}^2 (\tilde{Y}_\rho^1)_{[0]} \bar{\theta}^2 \bar{\theta}_{<0>}^2 (\tilde{X}_\lambda^3)_{<0> <0>} \tilde{x}_\rho^1) \\ (1.47) \quad &= (\tilde{x}_\lambda^1 \theta^1 \tilde{X}_\lambda^1 \cdot \varphi \cdot \theta_1^3 \bar{\theta}_1^3 \tilde{Y}_\rho^2 (\tilde{X}_\lambda^3)_{<0> <1>} \tilde{x}_\rho^2) \\ & \quad (\tilde{x}_\lambda^2 \theta_{[-1]}^2 \bar{\theta}^1 \tilde{X}_\lambda^2 \cdot \varphi' \cdot \theta_2^3 \bar{\theta}_2^3 \tilde{Y}_\rho^3 (\tilde{X}_\lambda^3)_{<1>} \tilde{x}_\rho^3 \natural \tilde{x}_\lambda^3 \theta_{[0]}^2 \bar{\theta}^2 \tilde{Y}_\rho^1 (\tilde{X}_\lambda^3)_{<0> <0>} \tilde{x}_\rho^1) \\ (1.46) \quad &= (\theta_1^1 \tilde{x}_\lambda^1 \tilde{X}_\lambda^1 \cdot \varphi \cdot \theta_1^3 (\tilde{x}_\lambda^3)_{<1>1} \tilde{Y}_\rho^2 (\tilde{X}_\lambda^3)_{<0> <1>} \tilde{x}_\rho^2) \\ & \quad (\theta_2^1 \tilde{x}_\lambda^2 \tilde{X}_\lambda^2 \cdot \varphi' \cdot \theta_2^3 (\tilde{x}_\lambda^3)_{<1>2} \tilde{Y}_\rho^3 (\tilde{X}_\lambda^3)_{<1>} \tilde{x}_\rho^3 \natural \theta^2 (\tilde{x}_\lambda^3)_{<0>} \tilde{Y}_\rho^1 (\tilde{X}_\lambda^3)_{<0> <0>} \tilde{x}_\rho^1) \\ (1.29) \quad &= (\theta_1^1 \cdot \varphi \cdot \theta_1^3) (\theta_2^1 \cdot \varphi' \cdot \theta_2^3) \natural \theta^2 \\ &= \Lambda(\varphi \varphi'), \end{aligned}$$

finishing the proof of (7.3). We can thus apply Proposition 7.2, obtaining an algebra map $w : \mathcal{A} \bowtie \mathbb{A} \rightarrow \mathcal{A} \natural \mathbb{A}$ satisfying $w \circ \Gamma = \Lambda$ and $w \circ j = j$, and given by the formula

$$\begin{aligned} w(\varphi \bowtie u) &= j(\tilde{q}_\rho^1) \Lambda(\varphi \cdot \tilde{q}_\rho^2) j(u) \\ &= (\tilde{q}_\rho^1)_{[-1]} \theta^1 \cdot \varphi \cdot \tilde{q}_\rho^2 \theta^3 u_{<1>} \natural (\tilde{q}_\rho^1)_{[0]} \theta^2 u_{<0>}. \end{aligned}$$

On the other hand, by using (1.47) and (1.5), one can easily check the following relation:

$$(\tilde{q}_\rho^1)_{[-1]} \theta^1 \otimes (\tilde{q}_\rho^1)_{[0]} \theta^2 \otimes \tilde{q}_\rho^2 \theta^3 = \Theta^1 \otimes \tilde{q}_\rho^1 \Theta_{<0>}^2 \otimes S^{-1}(\Theta^3) \tilde{q}_\rho^2 \Theta_{<1>}^2,$$

which implies that the map w above coincides with the algebra isomorphism ν given by (1.60).

As a consequence of Proposition 7.2, we immediately obtain the following new kind of universal property for the quasi-Hopf smash product:

Proposition 7.4 *Let H be a quasi-Hopf algebra and A a left H -module algebra. Denote by $i : A \rightarrow A \# H$, $i(a) = a \# 1$ and $j : H \rightarrow A \# H$, $j(h) = 1 \# h$. Let B be an associative algebra, $\gamma : H \rightarrow B$ an algebra map and $v : A \rightarrow B$ a linear map satisfying the following conditions:*

$$\gamma(h)v(a) = v(h_1 \cdot a)\gamma(h_2), \quad (7.8)$$

$$v(aa') = v(X^1 \cdot a)v(X^2 \cdot a')\gamma(X^3), \quad (7.9)$$

$$v(1_A) = 1_B, \quad (7.10)$$

for $a, a' \in A$, $h \in H$. Then there exists a unique algebra map $w : A \# H \rightarrow B$ such that $w \circ i = v$ and $w \circ j = \gamma$. Moreover, w is given by the formula $w(a \# h) = v(a)\gamma(h)$, for $a \in A$, $h \in H$.

The relation between the two universal properties of $A\#H$ is the following: Proposition 7.4 implies the usual universal property (unlike the case of ordinary Hopf algebras, the converse does not seem to hold). Indeed, if B is an associative algebra, $\gamma : H \rightarrow B$ an algebra map and $u : A \rightarrow B^\gamma$ a morphism of left H -module algebras, define the map

$$v : A \rightarrow B, \quad v(a) = \gamma(q^1)u(a)\gamma(S(q^2)),$$

where $q_R = q^1 \otimes q^2$ is given by (1.44). Then one can check that the maps γ and v satisfy the hypotheses of Proposition 7.4, and obviously the unique algebra map $w : A\#H \rightarrow B$ provided by Proposition 7.4 coincides with the map $u\# \gamma$ given by (1.25).

Proposition 7.4 may be easily extended to a universal property of the two-sided smash product:

Proposition 7.5 *Let H be a quasi-Hopf algebra, A a left H -module algebra and B a right H -module algebra. Denote by i_A, i_B, j the standard inclusions of A, B, H respectively into $A\#H\#B$. Let X be an associative algebra, $\gamma : H \rightarrow X$ an algebra map and $v_A : A \rightarrow X, v_B : B \rightarrow X$ two linear maps satisfying the following conditions:*

$$\begin{aligned} \gamma(h)v_A(a) &= v_A(h_1 \cdot a)\gamma(h_2), & v_A(aa') &= v_A(X^1 \cdot a)v_A(X^2 \cdot a')\gamma(X^3), \\ v_B(b)\gamma(h) &= \gamma(h_1)v_B(b \cdot h_2), & v_B(bb') &= \gamma(X^1)v_B(b \cdot X^2)v_B(b' \cdot X^3), \\ v_A(1_A) &= 1_X = v_B(1_B), & v_B(b)v_A(a) &= v_A(x^1 \cdot a)\gamma(x^2)v_B(b \cdot x^3), \end{aligned}$$

for all $a, a' \in A, b, b' \in B, h \in H$. Then there exists a unique algebra map $w : A\#H\#B \rightarrow X$ such that $w \circ i_A = v_A, w \circ i_B = v_B$ and $w \circ j = \gamma$. Moreover, w is given by the formula $w(a\#h\#b) = v_A(a)\gamma(h)v_B(b)$, for all $a \in A, h \in H, b \in B$.

For instance, the algebra isomorphism $A\#H\#B \simeq (A \otimes B) \sharp H$ from [28] may be easily reobtained by using this universal property.

References

- [1] S. E. Akrami and S. Majid, Braided cyclic cocycles and nonassociative geometry, *J. Math. Phys.* **45** (2004), 3883–3911.
- [2] H. Albuquerque, S. Majid, Quasialgebra structure of the octonions, *J. Algebra* **220** (1999), 188–224.
- [3] H. Albuquerque, S. Majid, Clifford algebras obtained by twisting of group algebras, *J. Pure Appl. Algebra* **171** (2002), 133–148.
- [4] J. Beck, Distributive laws, *Lecture Notes in Mathematics* **80** (1969), 119–140.
- [5] E. J. Beggs and S. Majid, Quantization by cochain twists and nonassociative differentials, arXiv:math.QA/0506450.
- [6] Y. Bespalov, B. Drabant, Cross product bialgebras. II, *J. Algebra* **240** (2001), 445–504.
- [7] P. Bieliavsky, P. Bonneau, Y. Maeda, Universal deformation formulae for three-dimensional solvable Lie groups, in “Quantum field theory and noncommutative geometry”, 127–141, *Lecture Notes in Phys.*, **662**, Springer, Berlin, 2005.

- [8] P. Bonneau, M. Gerstenhaber, A. Giaquinto, D. Sternheimer, Quantum groups and deformation quantization: Explicit approaches and implicit aspects, *J. Math. Phys.* **45** (2004), 3703–3741.
- [9] D. Bulacu and S. Caenepeel, Two-sided two-cosided Hopf modules and Doi-Hopf modules for quasi-Hopf algebras, *J. Algebra* **270** (2003), 55–95.
- [10] D. Bulacu, E. Nauwelaerts, Relative Hopf modules for (dual) quasi-Hopf algebras, *J. Algebra* **229** (2000), 632–659.
- [11] D. Bulacu, E. Nauwelaerts, Radford’s biproduct for quasi-Hopf algebras and bosonization, *J. Pure Appl. Algebra* **174** (2002), 1–42.
- [12] D. Bulacu, F. Panaite, F. Van Oystaeyen, Quasi-Hopf algebra actions and smash products, *Comm. Algebra* **28** (2000), 631–651.
- [13] D. Bulacu, F. Panaite, F. Van Oystaeyen, Generalized diagonal crossed products and smash products for quasi-Hopf algebras. Applications, *Comm. Math. Phys.* **266** (2006), 355–399.
- [14] S. Caenepeel, B. Ion, G. Militaru, S. Zhu, The factorization problem and the smash biproduct of algebras and coalgebras, *Algebr. Represent. Theory* **3** (2000), 19–42.
- [15] A. Cap, H. Schichl, J. Vanzura, On twisted tensor products of algebras, *Comm. Algebra* **23** (1995), 4701–4735.
- [16] V. G. Drinfeld, Quasi-Hopf algebras, *Leningrad Math. J.* **1** (1990), 1419–1457.
- [17] F. Hausser, F. Nill, Diagonal crossed products by duals of quasi-quantum groups, *Rev. Math. Phys.* **11** (1999), 553–629.
- [18] P. Jara Martínez, J. López Peña, F. Panaite, F. Van Oystaeyen, On iterated twisted tensor products of algebras, arXiv:math.QA/0511280.
- [19] C. Kassel, ”Quantum groups”, *Graduate Texts in Mathematics* **155**, Springer Verlag, Berlin, 1995.
- [20] J. López Peña, F. Panaite, F. Van Oystaeyen, General twisting of algebras, *Adv. Math.* (2006), doi:10.1016/j.aim.2006.10.003.
- [21] S. Majid, ”Foundations of quantum group theory”, Cambridge Univ. Press, 1995.
- [22] S. Majid, Quasi- $\ast$ structure on q -Poincaré algebras, *J. Geom. Phys.* **22** (1997), 14–58.
- [23] S. Majid, Quantum double for quasi-Hopf algebras, *Lett. Math. Phys.* **45** (1998), 1–9.
- [24] S. Majid, Gauge theory on nonassociative spaces, *J. Math. Phys.* **46** (2005), 103519, 23 pp.
- [25] M. Markl, Distributive laws and Koszulness, *Ann. Inst. Fourier* **46** (1996), 307–323.
- [26] S. Montgomery, ”Hopf algebras and their actions on rings”, CBMS Regional Conference Series, Vol. 82, Amer. Math. Soc., Providence, RI, 1993.
- [27] F. Panaite, F. Van Oystaeyen, Quasi-Hopf algebras and representations of octonions and other quasialgebras, *J. Math. Phys.* **45** (2004), 3912–3929.

- [28] F. Panaite, F. Van Oystaeyen, L-R-smash product for (quasi-) Hopf algebras, *J. Algebra* (2006), doi:10.1016/j.jalgebra.2006.07.020.
- [29] P. Schauenburg, Actions of monoidal categories and generalized Hopf smash products, *J. Algebra* **270** (2003), 521-563.
- [30] R. Street, The formal theory of monads, *J. Pure Appl. Algebra* **2** (1972), 149–168.
- [31] A. Van Daele, S. Van Keer, The Yang–Baxter and Pentagon equation, *Compositio Math.* **91** (1994), 201–221.
- [32] G. P. Wene, A construction relating Clifford algebras and Cayley-Dickson algebras, *J. Math. Phys.* **25** (1984), 2351–2353.